

Prof. S. Smith: Fall 1995

Problem 1: (a) Find the row-reduced echelon form of

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}.$$

$$\begin{array}{c} A_2^{-4 \times 1}, A_3^{-7 \times 1} \\ \rightarrow \end{array} \begin{pmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & -6 & -12 \end{pmatrix} \begin{array}{c} M_{-\frac{1}{3} \times 2} \\ \rightarrow \end{array} \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & -6 & -12 \end{pmatrix} \begin{array}{c} A_1^{-2 \times 2}, A_3^{6 \times 2} \\ \rightarrow \end{array} \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}$$

(b) What are the solutions of the system $Ax = 0$? (Check!)*Third variable is free, so solutions $x_3(1, -2, 1)^T$.***Problem 2:** Give the LU -decomposition of

$$A = \begin{pmatrix} 1 & 1 \\ 1 & 3 \end{pmatrix};$$

that is, find lower-triangular L and upper-triangular U , so that $A = LU$.*Get U from $A_2^{-1 \times 1} \rightarrow \begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix}$. So L from inverse operation $A_2^{+1 \times 1} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$* **Problem 3:** Use Cramer's rule (determinants) to solve $Ax = b$ given by

$$(A|b) = \left(\begin{array}{cc|c} 1 & 1 & 5 \\ 1 & 2 & 7 \end{array} \right).$$

*First $\det \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} = 1$, so $x_1 = \left(\frac{1}{1}\right) \det \begin{pmatrix} 5 & 1 \\ 7 & 2 \end{pmatrix} = 3$ and $x_2 = \det \begin{pmatrix} 1 & 5 \\ 1 & 7 \end{pmatrix} = 2$.***Problem 4:** (a) Find the inverse (by any method) of

$$A = \begin{pmatrix} 1 & 2 \\ 3 & 5 \end{pmatrix}.$$

Quick via adjoint: $\det \begin{pmatrix} 1 & 2 \\ 3 & 5 \end{pmatrix} = -1$, so inverse is $-\begin{pmatrix} 5 & -2 \\ -3 & 1 \end{pmatrix} = \begin{pmatrix} -5 & 2 \\ 3 & -1 \end{pmatrix}$.(b) Use the above to express the solutions of $Ax = b$ in terms of the constants b_1 and b_2 .*By $A^{-1}b$, namely $x_1 = -5b_1 + 2b_2$ and $x_2 = 3b_1 - b_2$.*

Problem 5: (a) Is $(1, 2, 3)$ in the span of $(4, 0, 5)$ and $(6, 0, 7)$?

No—for example, any linear combination of the latter two vectors has 0 in second entry.

(b) Let V be the space of all functions (with at least 2 derivatives).

Let W be the subset of all functions f which are solutions of the differential equation

$$f'' + 5f = 0.$$

Show that the solution set W is a subspace of V .

Take $f, g \in W$, and scalar c : Then we have $f'' + 5f = 0 = g'' + 5g$.

Is $f + g \in W$? $(f + g)'' + 5(f + g) = (f'' + 5f) + (g'' + 5g) = 0 + 0 = 0$, so OK.

Is $cf \in W$? $(cf)'' + 5(cf) = c(f'' + 5f) = c(0) = 0$, also OK.