

All 5 problems are worth 20 points each. You must SHOW WORK to receive credit.  
(If you use a calculator, WRITE "I used calculator" at those places).

**Problem 1:** Let  $A = \begin{pmatrix} 3 & 2 \\ 3 & 8 \end{pmatrix}$ .

(a) Find the characteristic polynomial, and the eigenvalues, of  $A$ .

$\det(A - xI) = (x - 3)(x - 8) - 6 = x^2 - 11x + 18 = (x - 9)(x - 2)$ , so eigenvalues are 2, 9.

(b) Find the eigenspaces for those eigenvalues.

For 2:  $A - 2I = \begin{pmatrix} 1 & 2 \\ 3 & 6 \end{pmatrix}$ , via  $A_2^{-2 \times 1}$  to  $\begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix}$ , get solutions  $a(-2, 1)^T$ .

For 9:  $A - 9I = \begin{pmatrix} -6 & 2 \\ 3 & -1 \end{pmatrix}$ , via  $A_2^{\frac{1}{2} \times 1}$  to  $\begin{pmatrix} -6 & 2 \\ 0 & 0 \end{pmatrix}$ , get solutions  $b(\frac{1}{3}, 1)^T$ .

**Problem 2:** Given the differential equation system (functions of  $t$ ):  $\begin{pmatrix} y_1' = 2y_1 + 3y_2 \\ y_2' = -y_1 - 2y_2 \end{pmatrix}$ .

I GIVE you the information that eigenvalues of the coefficient matrix  $A$  for this system are 1, -1,  
(a) Find eigenvectors for  $A$ ; then use them to give the *general* solution of the system (with undetermined constants  $c_1, c_2$ ).

For 1, get  $a(-3, 1)^T$ ; for -1, get  $b(-1, 1)^T$ .

Then solution vector  $c_1 \begin{pmatrix} -3 \\ 1 \end{pmatrix} e^t + c_2 \begin{pmatrix} -1 \\ 1 \end{pmatrix} e^{-t}$  so  $y_1 = -3c_1 e^t - c_2 e^{-t}$  and  $y_2 = c_1 e^t + c_2 e^{-t}$ .

(b) Now find the particular solution (values of  $c_1, c_2$ ) given initial values  $y_1(0) = 3, y_2(0) = 2$ .

Solve  $\left( \begin{array}{cc|c} -3 & -1 & 3 \\ 1 & 1 & 2 \end{array} \right)$  to get  $c_1 = -2.5, c_2 = 4.5$ .

So  $y_1 = 7.5e^t - 4.5e^{-t}$  and  $y_2 = -2.5e^t + 4.5e^{-t}$ .

**Problem 3:** (a) Let  $A = \begin{pmatrix} 4 & 0 & 0 \\ 0 & 5 & 0 \\ 1 & 0 & 4 \end{pmatrix}$ .

Is  $A$  diagonalizable? Indicate why/why not.

No: As  $A$  is triangular, we see the eigenvalues are 4,4,5. Solving  $(A - 4I)x = 0$ , we see the eigenspace for 4 consists of  $a(0, 0, 1)^T$ : dimension only 1, whereas the eigenvalue is repeated twice. (That is, the geometric multiplicity is less than the algebraic multiplicity). So we cannot get a basis of eigenvectors, and  $A$  is not diagonalizable.

(b) For  $A = \begin{pmatrix} 0 & 2 \\ 2 & 0 \end{pmatrix}$ , I GIVE you that eigenvalues are 2, -2; with corresponding eigenvectors  $(1, 1)^T$  and  $(1, -1)^T$ . Use the diagonalization of  $A$  (namely use the relevant matrix  $X$  with  $X^{-1}AX$  diagonal) to determine the 5-th power  $A^5$ .

We can use  $X = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$ ,  $X^{-1} = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$  with  $D = X^{-1}AX = \begin{pmatrix} 2 & 0 \\ 0 & -2 \end{pmatrix}$ .

So  $A = XDX^{-1}$  and then  $A^5 = X(D^5)X^{-1}$  where  $D^5 = \begin{pmatrix} 2^5 & 0 \\ 0 & (-2)^5 \end{pmatrix}$ .

Multiplying out we get  $A^5 = \begin{pmatrix} 0 & 32 \\ 32 & 0 \end{pmatrix}$ .

**Problem 4:** Let  $A$  be the symmetric matrix  $\begin{pmatrix} 5 & -4 & -2 \\ -2 & 2 & 2 \end{pmatrix}$

with eigenvalues  $1, 1, 10$  of  $A$ ; and an eigenvector  $(-2, 2, 1)^T$  for eigenvalue  $10$ .

Find the eigenspace of  $A$  for eigenvalue  $1$ .

The Jordan form of  $A - 1 \cdot I$  has  $(1, -1, -.5)$  as its only nonzero row.

Find a basis for the eigenspace  $\frac{1}{2}(c, b, c)^T$ ; and one possible basis is  $(1, 1, 0)^T$  and  $(\frac{1}{2}, 0, 1)^T$ .

(b) Give an orthogonal diagonalization of  $A$ ; that is, find an orthogonal matrix  $X$  (satisfying  $X^{-1} = X^T$ ) with  $X^{-1}AX$  is diagonal.

For  $10$ : space is 1-dimensional; divide original  $(-2, 2, 1)$  by its length  $3$ :  $x_3 = \frac{1}{3}(-2, 2, 1)^T$ .

For  $1$ : Start with above basis like  $v_1 = (1, 1, 0)^T$  and  $v_2 = (1, 0, 2)^T$ .

Apply Gram-Schmidt: first  $q_1 = (1, 1, 0)$

and then  $q_2 = v_2 - \frac{v_2 \cdot q_1}{q_1 \cdot q_1} q_1 = (1, 0, 2)^T - \frac{1}{2}(1, 1, 0)^T = (\frac{1}{2}, -\frac{1}{2}, 2)^T$

so may as well use the more convenient multiple  $q_2 = (1, -1, 4)^T$ .

Now divide each by length, to get  $x_1 = \frac{1}{\sqrt{2}}(1, 1, 0)^T$  and  $x_2 = \frac{1}{\sqrt{18}}(1, -1, 4)^T$ .

So can use  $X = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{18}} & -\frac{2}{3} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{18}} & \frac{2}{3} \\ 0 & \frac{4}{\sqrt{18}} & \frac{1}{3} \end{pmatrix}$ .

**Problem 5:** (a) For the Markov matrix  $A = \begin{pmatrix} .1 & .6 \\ .9 & .4 \end{pmatrix}$ , find the eigenvalues; also find the

“steady-state” vector  $v$ . (That is,  $Av = v$ , and the coordinates of  $v$  add up to  $1$ .)

$\det(A - xI) = (.1 - x)(.4 - x) - (.9)(.6) = x^2 - .5x + .04 - .54 = x^2 - .5x - .5 = (x - 1)(x + .5)$   
so the eigenvalues are  $1, -.5$ .

The eigenspace for  $1$  consists of vectors  $a(2, 3)^T$ . So the steady-state vector is  $(.4, .6)^T$ .

(b) Write the matrix  $\begin{pmatrix} 9 & -3 \\ -3 & 2 \end{pmatrix}$  as a product  $LDL^T$ , with  $L$  lower triangular, and  $D$  diagonal.

First get  $LU$ -decomposition as in Section 1.4:

$A_2^{\frac{1}{3} \times 1}$  takes  $A$  to  $U = \begin{pmatrix} 9 & -3 \\ 0 & 1 \end{pmatrix}$

so that  $L = \begin{pmatrix} 1 & 0 \\ -\frac{1}{3} & 1 \end{pmatrix}$

then factor  $U = DU^*$  by using diagonal values of  $U$  in  $D$ :

$D = \begin{pmatrix} 9 & 0 \\ 0 & 1 \end{pmatrix}$  so that  $U^* = \begin{pmatrix} 1 & -\frac{1}{3} \\ 0 & 1 \end{pmatrix}$ , and then observe that  $U^*$  is indeed  $L^T$ .