

Prof. S. Smith (originally Prof. H. Alexander): Tues 4 May 1999

There are 6 problems, worth 20 % each (so you can get more than 100 %).

You must SHOW WORK to get credit; if by calculator, show WHERE and HOW you used it.

**Problem 1:** (a) (as promised, a flashback to the midterm):

Let  $S$  be the set of all lower triangular  $2 \times 2$  matrices. Remember that these matrices have the general form  $\begin{pmatrix} a & 0 \\ b & c \end{pmatrix}$ . Show that  $S$  is a subspace of the vector space  $\mathbf{R}^{2 \times 2}$  of all  $2 \times 2$  matrices.

$$(\text{closure, } +: ) \begin{pmatrix} a & 0 \\ b & c \end{pmatrix} + \begin{pmatrix} d & 0 \\ e & f \end{pmatrix} = \begin{pmatrix} a+d & 0 \\ b+e & c+f \end{pmatrix}, \text{ also lower-triangular, so in } S.$$

$$(\text{closure, scalar mult:}) f \begin{pmatrix} a & 0 \\ b & c \end{pmatrix} = \begin{pmatrix} fa & 0 \\ fb & fc \end{pmatrix}, \text{ also lower-triangular, so in } S.$$

(b) Give the standard matrix representation for linear transformation from  $\mathbf{R}^3$  to  $\mathbf{R}^2$  defined by  $L[(x_1, x_2, x_3)^T] = (x_1 - 3x_2 + 2x_3, 7x_2 + 3x_3)^T$ . (That is, just use the standard basis for each space).

$$\text{Apply } L \text{ to each basis vector, and write as columns: } \begin{pmatrix} 1 & -3 & 2 \\ 0 & 7 & 3 \end{pmatrix}$$

**Problem 2:** (a) Find the orthogonal complement  $S^\perp$  to the space spanned by the two vectors  $(1, 1, 1)^T$  and  $(1, 0, 2)^T$ .

$$\text{Write as rows of } A: \begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 2 \end{pmatrix}. \text{ Then } Ax = 0 \text{ has solutions } x_3(-2, 1, 1)^T.$$

(b) Find the least-squares solution  $\hat{x}$  of the inconsistent system  $Ax = b$  given by:  $\left( \begin{array}{cc|c} 0 & 1 & 1 \\ 1 & 1 & 2 \\ 2 & 1 & 4 \end{array} \right)$ .

Give the projection  $p$  of  $b$  in the column space of  $A$ . What is the error?

$$\text{Multiply by } A^T = \begin{pmatrix} 0 & 1 & 2 \\ 1 & 1 & 1 \end{pmatrix} \text{ to get normal equations } \left( \begin{array}{cc|c} 5 & 3 & 10 \\ 3 & 3 & 7 \end{array} \right).$$

$$\text{Solve to get } \hat{x} = \frac{1}{6}(9, 5)^T. \text{ Then } p = A\hat{x} = \frac{1}{6}(5, 14, 23)^T. \text{ And error} = b - p = \frac{1}{6}(1, -2, 1)^T.$$

**Problem 3:** (a) Use the Gram-Schmidt process to find an orthonormal basis for the space spanned by the two vectors  $b_1 = (0, 1, 2)^T$  and  $b_2 = (1, 1, 1)^T$ .

$$(\text{Working with rows}) x_1 = (0, 1, 2) \text{ and } x_2 = b_2 - \frac{x_2 \cdot x_1}{x_1 \cdot x_1} x_1 = (1, 1, 1) - \frac{3}{5}(0, 1, 2) = (1, \frac{2}{5}, -\frac{1}{5})$$

$$\text{so } u_1 = x_1/|x_1| = \frac{1}{\sqrt{5}}(0, 1, 2) \text{ and } u_2 = \frac{1}{\sqrt{30}}(5, 2, -1).$$

(b) Find the projection  $p$  of the vector  $b = (1, 2, 3)^T$  in the 2-dimensional subspace spanned by the vectors  $b_1 = (1, -1, 0)^T$  and  $b_2 = (1, 1, 1)^T$ .

Many ways of solving. Easiest: note  $b_1$  and  $b_2$  are orthogonal.

So  $p$  has coordinates  $(b \cdot b_1)/(b_1 \cdot b_1) = -\frac{1}{2}$  and similarly for  $b_2$  to get 2.

$$\text{so } p = -\frac{1}{2}(1, -1, 0)^T + 2(1, 1, 1)^T = \frac{1}{2}(3, 5, 4)^T.$$

**Problem 4:** For the matrix  $A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}$ :

(a) Find the eigenvalues of  $A$ .

$\det(A - xI) = x^2 - 5x + 4 - 4 = x^2 - 5x = x(x - 5)$ , so eigenvalues are 0, 5.

(b) For each eigenvalue, determine the corresponding eigenvectors.

( $\lambda = 0$ ) For  $A - 0I = A$  solve  $Ax = 0$ : solutions are multiples of  $(-2, 1)^T$ .

( $\lambda = 5$ ) solve  $(A - 5I)x = 0$  where  $A - 5I = \begin{pmatrix} -4 & 2 \\ 2 & -1 \end{pmatrix}$  solutions are multiples of  $(1, 2)^T$ .

**Problem 5:** Given the differential equation system ( $y$  as functions of  $t$ ):  $\begin{pmatrix} y_1' = y_1 + 2y_2 \\ y_2' = 2y_1 + y_2 \end{pmatrix}$ .

I GIVE you the information that eigenvalues are 3,  $-1$ ; with eigenvectors are  $(1, 1)^T$  and  $(1, -1)^T$ .

(a) Give the *general* solution of the system (with undetermined constants  $c_1, c_2$ ).

$\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} c_1 e^{3t} \\ c_2 e^{-t} \end{pmatrix}$ , so  $y_1 = c_1 e^{3t} + c_2 e^{-t}$  and  $y_2 = c_1 e^{3t} - c_2 e^{-t}$ .

(b) Now determine the values of  $c_1, c_2$  for the initial value problem  $y_1(0) = 4, y_2(0) = 2$ .

Solve  $\begin{pmatrix} 1 & 1 & | & 4 \\ 1 & -1 & | & 2 \end{pmatrix}$  to get  $c_1 = 3, c_2 = 1$ . So  $y_1 = 3e^{3t} + e^{-t}$  and  $y_2 = 3e^{3t} - e^{-t}$ .

**Problem 6:** Let  $A$  be the matrix  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ . I GIVE you that the eigenvalues of  $A$  are  $-1, 1$ .

(a) Diagonalize  $A$ : that is, find the eigenvalues and eigenvectors, and give a matrix  $X$  such that  $X^{-1}AX$  is a diagonal matrix  $D$ .

For 1: eigenvectors are  $a(1, 1)^T$ , For  $-1$ : Get eigenvectors  $b(1, -1)^T$ . So  $X = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$ .

(b) Now let  $A$  be the symmetric matrix  $\begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}$ .

I GIVE you that the eigenvalues of  $A$  are  $-1, -1, 2$ ; and that corresponding eigenvectors are  $(1, -1, 0)^T$ ,  $(1, 0, -1)^T$ , and  $(1, 1, 1)^T$ . Give an *orthogonal* diagonalization of  $A$ :

That is, use Gram-Schmidt to find an orthonormal basis for each eigenspace. Then use that to build an orthogonal matrix  $X$  (that is,  $X^{-1} = X^T$ ) with  $X^{-1}AX$  diagonal.

Apply Gram-Schmidt to the  $-1$ -eigenspace to get  $\frac{1}{\sqrt{2}}(1, -1, 0)^T$  and  $\frac{1}{\sqrt{6}}(1, 1, -2)^T$ .

For the 2-eigenspace, use  $\frac{1}{\sqrt{3}}(1, 1, 1)^T$ .

So can use  $X = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ 0 & -\frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} \end{pmatrix}$ .