

Math 181, Exam 1, Spring 2009
Problem 1 Solution

1. Differentiate the function:

$$F(x) = \int_{\sqrt{x}}^{x^2} e^{t^3} dt$$

Solution: Using the Fundamental Theorem of Calculus Part II and the Chain Rule, the derivative of $F(x) = \int_{g(x)}^{h(x)} f(t) dt$ is:

$$\begin{aligned} F'(x) &= \frac{d}{dx} \int_{g(x)}^{h(x)} f(t) dt \\ &= f(h(x)) \cdot \frac{d}{dx} h(x) - f(g(x)) \cdot \frac{d}{dx} g(x) \end{aligned}$$

Applying the formula to the given function $F(x)$ we get:

$$\begin{aligned} F'(x) &= \frac{d}{dx} \int_{\sqrt{x}}^{x^2} e^{t^3} dt \\ &= e^{(x^2)^3} \cdot \frac{d}{dx} (x^2) - e^{(\sqrt{x})^3} \cdot \frac{d}{dx} (\sqrt{x}) \\ &= \boxed{e^{x^6} \cdot (2x) - e^{x^{3/2}} \cdot \left(\frac{1}{2\sqrt{x}}\right)} \end{aligned}$$

Math 181, Final Exam, Study Guide
Problem 2 Solution

2. Compute the definite integral:

$$\int_0^1 x e^{3x} dx$$

Solution: We will evaluate the integral using Integration by Parts. Let $u = x$ and $v' = e^{3x}$. Then $u' = 1$ and $v = \frac{1}{3}e^{3x}$. Using the Integration by Parts formula:

$$\int_a^b uv' dx = [uv]_a^b - \int_a^b u'v dx$$

we get:

$$\begin{aligned} \int_0^1 x e^{3x} dx &= \left[\frac{1}{3} x e^{3x} \right]_0^1 - \int_0^1 \frac{1}{3} e^{3x} dx \\ &= \left[\frac{1}{3} x e^{3x} \right]_0^1 - \left[\frac{1}{9} e^{3x} \right]_0^1 \\ &= \left[\frac{1}{3} x e^{3x} - \frac{1}{9} e^{3x} \right]_0^1 \\ &= \left[\frac{1}{3}(1)e^{3(1)} - \frac{1}{9}e^{3(1)} \right] - \left[\frac{1}{3}(0)e^{3(0)} - \frac{1}{9}e^{3(0)} \right] \\ &= \boxed{\frac{2}{9}e^3 + \frac{1}{9}} \end{aligned}$$

Math 181, Exam 2, Study Guide
Problem 1 Solution

1. Compute the integrals:

$$\int \frac{x}{\sqrt{x-2}} dx \quad \int x^3 \sin(x^2) dx \quad \int \frac{dx}{x^2+x-6} dx$$

$$\int \frac{dx}{x^2+x+3} \quad \int \frac{dx}{x^3-x} \quad \int x^6 \ln x dx$$

$$\int \arctan x dx \quad \int \cos(\sqrt{x}) dx \quad \int x^2 e^{2x} dx$$

Solution:

- (1) The first integral is computed using the u -substitution method. Let $u = x - 2$. Then $du = dx$ and $x = u + 2$. Substituting these into the integral and evaluating we get:

$$\begin{aligned} \int \frac{x}{\sqrt{x-2}} dx &= \int \frac{u+2}{\sqrt{u}} du \\ &= \int (u^{1/2} + 2u^{-1/2}) du \\ &= \frac{2}{3}u^{3/2} + 4u^{1/2} + C \\ &= \boxed{\frac{2}{3}(x-2)^{3/2} + 4(x-2)^{1/2} + C} \end{aligned}$$

- (2) The second integral is computed using the u -substitution method. Let $u = x^2$. Then $du = 2x dx \Rightarrow \frac{1}{2} du = x dx$ and we get:

$$\begin{aligned} \int x^3 \sin(x^2) dx &= \int x^2 \sin(x^2) x dx \\ &= \int u \sin u \left(\frac{1}{2} du\right) \\ &= \frac{1}{2} \int u \sin u du \end{aligned}$$

We now use Integration by Parts to evaluate the above integral. Let $w = u$ and $v' = \sin u$. Then $w' = 1$ and $v = -\cos u$. Using the Integration by Parts formula:

$$\int wv' du = wv - \int w'v du$$

we get:

$$\begin{aligned}\int u \sin u \, du &= u(-\cos u) - \int 1 \cdot (-\cos u) \, du \\ &= -u \cos u + \int \cos u \, du \\ &= -u \cos u + \sin u + C\end{aligned}$$

Therefore,

$$\begin{aligned}\int x^3 \sin(x^2) \, dx &= \frac{1}{2} \int u \sin u \, du \\ &= \frac{1}{2} (-u \cos u + \sin u) + C \\ &= -\frac{1}{2} u \cos u + \frac{1}{2} \sin u + C \\ &= \boxed{-\frac{1}{2} x^2 \cos(x^2) + \frac{1}{2} \sin(x^2) + C}\end{aligned}$$

- (3) The third integral is computed using Partial Fraction Decomposition. Factoring the denominator and decomposing we get:

$$\frac{1}{x^2 + x - 6} = \frac{1}{(x+3)(x-2)} = \frac{A}{x+3} + \frac{B}{x-2}$$

Multiplying the equation by $(x+3)(x-2)$ we get:

$$1 = A(x-2) + B(x+3)$$

Next we plug in two different values of x to get a system of two equations in two unknowns (A , B). Letting $x = -3$ and $x = 2$ we get:

$$\begin{aligned}x = -3: \quad 1 &= A(-3-2) + B(-3+3) \quad \Rightarrow \quad A = -\frac{1}{5} \\ x = 2: \quad 1 &= A(2-2) + B(2+3) \quad \Rightarrow \quad B = \frac{1}{5}\end{aligned}$$

Plugging these values of A and B back into the decomposed equation and integrating we get:

$$\begin{aligned}\int \frac{1}{x^2 + x - 6} \, dx &= \int \left(\frac{-\frac{1}{5}}{x+3} + \frac{\frac{1}{5}}{x-2} \right) \, dx \\ &= \boxed{-\frac{1}{5} \ln|x+3| + \frac{1}{5} \ln|x-2| + C}\end{aligned}$$

(4) We begin solving the fourth integral by completing the square in the denominator.

$$\int \frac{1}{x^2 + x + 3} dx = \int \frac{1}{(x + \frac{1}{2})^2 + \frac{11}{4}} dx$$

We then evaluate the integral using the u -substitution method. Let $\frac{\sqrt{11}}{2} u = x + \frac{1}{2}$. Then $\frac{\sqrt{11}}{2} du = dx$ and we get:

$$\begin{aligned} \int \frac{1}{x^2 + x + 3} dx &= \int \frac{1}{(x + \frac{1}{2})^2 + \frac{11}{4}} dx \\ &= \frac{\sqrt{11}}{2} \int \frac{1}{\frac{11}{4}u^2 + \frac{11}{4}} du \\ &= \frac{2}{\sqrt{11}} \int \frac{1}{u^2 + 1} du \\ &= \frac{2}{\sqrt{11}} \arctan u + C \\ &= \boxed{\frac{2}{\sqrt{11}} \arctan \left[\frac{2}{\sqrt{11}} \left(x + \frac{1}{2} \right) \right] + C} \end{aligned}$$

(5) We will evaluate the fifth integral using Partial Fraction Decomposition. First, we factor the denominator and then decompose the rational function into a sum of simpler rational functions.

$$\frac{1}{x^3 - x} = \frac{1}{x(x-1)(x+1)} = \frac{A}{x} + \frac{B}{x-1} + \frac{C}{x+1}$$

Next, we multiply the above equation by $x(x-1)(x+1)$ to get:

$$1 = A(x-1)(x+1) + Bx(x+1) + Cx(x-1)$$

Then we plug in three different values for x to create a system of three equations in three unknowns (A, B, C) . We select $x = 0$, $x = -1$, and $x = 1$ for simplicity.

$$\begin{aligned} x = 0 : A(0-1)(0+1) = 1 &\Rightarrow -A = 1 \Rightarrow A = -1 \\ x = -1 : C(-1)(-1-1) = 1 &\Rightarrow 2C = 1 \Rightarrow C = \frac{1}{2} \\ x = 1 : B(1)(1+1) = 1 &\Rightarrow 2B = 1 \Rightarrow B = \frac{1}{2} \end{aligned}$$

Finally, we plug these values for A , B , and C back into the decomposition and integrate.

$$\begin{aligned} \int \frac{dx}{x^3 - 4x} &= \int \left(\frac{A}{x} + \frac{B}{x-1} + \frac{C}{x+1} \right) dx \\ &= \int \left(\frac{-1}{x} + \frac{\frac{1}{2}}{x-1} + \frac{\frac{1}{2}}{x+1} \right) dx \\ &= \boxed{-\ln|x| + \frac{1}{2}\ln|x-1| + \frac{1}{2}\ln|x+1| + C} \end{aligned}$$

- (6) The sixth integral is computed using Integration by Parts. Let $u = \ln x$ and $v' = x^6$. Then $u' = \frac{1}{x}$ and $v = \frac{1}{7}x^7$. Using the Integration by Parts formula:

$$\int uv' dx = uv - \int u'v dx$$

we get:

$$\begin{aligned} \int x^6 \ln x dx &= (\ln x) \left(\frac{1}{7}x^7 \right) - \int \frac{1}{x} \cdot \frac{1}{7}x^7 dx \\ &= \frac{1}{7}x^7 \ln x - \frac{1}{7} \int x^6 dx \\ &= \boxed{\frac{1}{7}x^7 \ln x - \frac{1}{49}x^7 + C} \end{aligned}$$

- (7) We will evaluate the seventh integral using Integration by Parts. Let $u = \arctan x$ and $v' = 1$. Then $u' = \frac{1}{x^2 + 1}$ and $v = x$. Using the Integration by Parts formula:

$$\int uv' dx = uv - \int u'v dx$$

we get:

$$\int \arctan x dx = x \arctan x - \int \frac{1}{x^2 + 1} x dx.$$

Use the substitution $w = x^2 + 1$ to evaluate the integral on the right hand side. Then $dw = 2x dx \Rightarrow \frac{1}{2}dw = x dx$ and we get:

$$\begin{aligned} \int \arctan x dx &= x \arctan x - \frac{1}{2} \int \frac{1}{w} dw \\ &= x \arctan x - \frac{1}{2} \ln|w| + C \\ &= \boxed{x \arctan x - \frac{1}{2} \ln(x^2 + 1) + C} \end{aligned}$$

Note that the absolute value signs aren't needed because $x^2 + 1 > 0$ for all x .

(8) To begin the solution of the eighth integral, we first use the u -substitution method.

Let $u = \sqrt{x}$. Then $du = \frac{1}{2\sqrt{x}} dx \Rightarrow 2u du = dx$ and we get:

$$\begin{aligned}\int \cos(\sqrt{x}) dx &= \int \cos u (2u du) \\ &= 2 \int u \cos u du\end{aligned}$$

We now use Integration by Parts to evaluate the above integral. Let $w = u$ and $v' = \cos u$. Then $w' = 1$ and $v = \sin u$. Using the Integration by Parts formula:

$$\int wv' du = wv - \int w'v du$$

we get:

$$\begin{aligned}\int u \cos u du &= u \sin u - \int \sin u du \\ \int u \cos u du &= u \sin u + \cos u + C\end{aligned}$$

Therefore,

$$\begin{aligned}\int \cos(\sqrt{x}) dx &= 2 \int u \cos u du \\ &= 2(u \sin u + \cos u) + C \\ &= 2u \sin u + \frac{1}{2} \cos u + C \\ &= \boxed{2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x}) + C}\end{aligned}$$

(9) The last integral is computed using Integration by Parts. Let $u = x^2$ and $v' = e^{2x}$. Then $u' = 2x$ and $v = \frac{1}{2}e^{2x}$. Using the Integration by Parts formula:

$$\int uv' dx = uv - \int u'v dx$$

we get:

$$\begin{aligned}\int x^2 e^{2x} dx &= \frac{1}{2}x^2 e^{2x} - \int 2x \left(\frac{1}{2}e^{2x}\right) dx \\ &= \frac{1}{2}x^2 e^{2x} - \int x e^{2x} dx\end{aligned}$$

A second Integration by Parts must be performed. Let $u = x$ and $v' = e^{2x}$. Then $u' = 1$ and $v = \frac{1}{2}e^{2x}$. Using the Integration by Parts formula again we get:

$$\begin{aligned}\int x^2 e^{2x} dx &= \frac{1}{2}x^2 e^{2x} - \left[\frac{1}{2}x e^{2x} - \frac{1}{2} \int e^{2x} dx \right] \\ &= \boxed{\frac{1}{2}x^2 e^{2x} - \frac{1}{2}x e^{2x} + \frac{1}{4}e^{2x} + C}\end{aligned}$$

Math 181, Final Exam, Study Guide
Problem 4 Solution

4. Determine if the following improper integrals converge or not.

$$\int_0^{+\infty} x e^{-2x} dx \quad \int_0^{+\infty} \frac{dx}{x^2 + 4} \quad \int_0^1 \frac{e^x}{\sqrt{x}} dx \quad \int_1^{+\infty} \frac{x^{3/2} + 3}{\sqrt{x}} dx$$

Solution: We evaluate the first integral by turning it into a limit calculation.

$$\int_0^{+\infty} x e^{-2x} dx = \lim_{R \rightarrow +\infty} \int_0^R x e^{-2x} dx$$

We use Integration by Parts to compute the integral. Let $u = x$ and $v' = e^{-2x}$. Then $u' = 1$ and $v = -\frac{1}{2}e^{-2x}$. Using the Integration by Parts formula we get:

$$\begin{aligned} \int_a^b uv' dx &= [uv]_a^b - \int_a^b u'v dx \\ \int_0^R x e^{-2x} dx &= \left[-\frac{1}{2} x e^{-2x} \right]_0^R - \int_0^R \left(-\frac{1}{2} e^{-2x} \right) dx \\ &= \left[-\frac{1}{2} x e^{-2x} \right]_0^R + \frac{1}{2} \int_0^R e^{-2x} dx \\ &= \left[-\frac{1}{2} x e^{-2x} \right]_0^R + \frac{1}{2} \left[-\frac{1}{2} e^{-2x} \right]_0^R \\ &= \left[-\frac{1}{2} R e^{-2R} + \frac{1}{2} (0) e^{-2(0)} \right] + \frac{1}{2} \left[-\frac{1}{2} e^{-2R} + \frac{1}{2} e^{-3(0)} \right] \\ &= -\frac{R}{2e^{2R}} - \frac{1}{4e^{2R}} + \frac{1}{4} \end{aligned}$$

We now take the limit of the above function as $R \rightarrow +\infty$.

$$\begin{aligned} \int_0^{+\infty} x e^{-2x} dx &= \lim_{R \rightarrow +\infty} \int_0^R x e^{-2x} dx \\ &= \lim_{R \rightarrow +\infty} \left(-\frac{R}{2e^{2R}} - \frac{1}{4e^{2R}} + \frac{1}{4} \right) \\ &= -0 - 0 + \frac{1}{4} \\ &= \boxed{\frac{1}{4}} \end{aligned}$$

We evaluate the second integral as follows:

$$\begin{aligned}
 \int_0^{+\infty} \frac{dx}{x^2 + 4} &= \lim_{R \rightarrow +\infty} \int_0^R \frac{dx}{x^2 + 4} \\
 &= \lim_{R \rightarrow +\infty} \left[\frac{1}{2} \arctan \left(\frac{x}{2} \right) \right]_0^R \\
 &= \lim_{R \rightarrow +\infty} \left[\frac{1}{2} \arctan \left(\frac{R}{2} \right) - \frac{1}{2} \arctan \left(\frac{0}{2} \right) \right] \\
 &= \frac{1}{2} \left(\frac{\pi}{2} \right) - \frac{1}{2} (0) \\
 &= \boxed{\frac{\pi}{4}}
 \end{aligned}$$

We will use the Comparison Test to show that the third integral converges. Let $g(x) = \frac{e^x}{\sqrt{x}}$. We must choose a function $f(x)$ that satisfies:

$$(1) \int_0^1 f(x) dx \text{ converges} \quad \text{and} \quad (2) \quad 0 \leq g(x) \leq f(x) \text{ for } 0 \leq x \leq 1$$

We choose $f(x) = \frac{e}{\sqrt{x}}$. This function satisfies the inequality:

$$\begin{aligned}
 0 &\leq g(x) \leq f(x) \\
 0 &\leq \frac{e^x}{\sqrt{x}} \leq \frac{e}{\sqrt{x}}
 \end{aligned}$$

for $0 \leq x \leq 1$ because the denominator of $g(x)$ is greater than the denominator of $f(x)$ for these values of x . Furthermore, the integral $\int_0^1 f(x) dx = \int_0^1 \frac{e}{\sqrt{x}} dx$ converges because it is a p -integral with $p = \frac{1}{2} < 1$. Therefore, the integral $\int_0^1 g(x) dx = \int_0^1 \frac{e^x}{\sqrt{x}} dx$ converges by the Comparison Test.

We split the fourth integral into two integrals.

$$\int_1^{+\infty} \frac{x^{3/2} + 3}{\sqrt{x}} dx = \int_1^{+\infty} \frac{x^{3/2}}{\sqrt{x}} dx + \int_1^{+\infty} \frac{3}{\sqrt{x}} dx = \int_1^{+\infty} x dx + 3 \int_1^{+\infty} \frac{dx}{\sqrt{x}}$$

Both integrals on the far right hand side of the above equation are divergent p -integrals ($p = -1$ and $p = \frac{1}{2}$, both of which are less than 1). Thus, the integral diverges.

Math 181, Final Exam, Study Guide
Problem 5 Solution

5. Let R be the region included by the curves $y = 0$ and $y = x^2 + x$ between $x = 0$ and $x = 1$. Find the volume of the solid of revolution when the axis is the line $y = -1$.

Solution: We find the volume using the Washer method. The formula we will use is:

$$V = \pi \int_a^b [(\text{top} - (-1))^2 - (\text{bottom} - (-1))^2] dx$$

where the top curve is $y = x + x^2$ and the bottom curve is $y = 0$. The lower limit is $a = 0$ and the upper limit is $b = 1$. The volume is then:

$$\begin{aligned} V &= \pi \int_a^b [(\text{top} - (-1))^2 - (\text{bottom} - (-1))^2] dx \\ &= \pi \int_0^1 [(x + x^2 + 1)^2 - (0 + 1)^2] dx \\ &= \pi \int_0^1 (x^2 + x^3 + x + x^3 + x^4 + x^2 + x + x^2 + 1 - 1) dx \\ &= \pi \int_0^1 (x^4 + 2x^3 + 3x^2 + 2x) dx \\ &= \pi \left[\frac{x^5}{5} + \frac{x^4}{2} + x^3 + x^2 \right]_0^1 \\ &= \pi \left[\frac{1}{5} + \frac{1}{2} + 1 + 1 \right] \\ &= \boxed{\frac{27\pi}{10}} \end{aligned}$$

Math 181, Final Exam, Study Guide
Problem 6 Solution

6. Find the arclength of the graph of the function $y = 2x^{3/2} + 5$ between $x = 0$ and $x = 1$.

Solution: The arclength is:

$$\begin{aligned} L &= \int_a^b \sqrt{1 + (y')^2} dx \\ &= \int_0^1 \sqrt{1 + (3x^{1/2})^2} dx \\ &= \int_0^1 \sqrt{1 + 9x} dx \end{aligned}$$

We now use the u -substitution $u = 1 + 9x$. Then $\frac{1}{9} du = dx$, the lower limit of integration changes from 0 to 1, and the upper limit of integration changes from 1 to 10.

$$\begin{aligned} L &= \int_0^2 \sqrt{1 + 9x} dx \\ &= \frac{1}{9} \int_1^{10} \sqrt{u} du \\ &= \frac{1}{9} \left[\frac{2}{3} u^{3/2} \right]_1^{10} \\ &= \frac{1}{9} \left[\frac{2}{3} (10)^{3/2} - \frac{2}{3} (1)^{3/2} \right] \\ &= \boxed{\frac{2}{27} [10\sqrt{10} - 1]} \end{aligned}$$

Math 181, Final Exam, Study Guide
Problem 7 Solution

7. Let $f(x) = x^2$ on the interval $[0, 1]$. Compute Mid(3) and Trap(3). Which one is an overestimate and why?

Solution: The length of each subinterval of $[0, 1]$ is:

$$\Delta x = \frac{b - a}{N} = \frac{1 - 0}{3} = \frac{1}{3}$$

The estimate Mid(3) is:

$$\begin{aligned} \text{Mid}(3) &= \Delta x \left[f\left(\frac{1}{6}\right) + f\left(\frac{3}{6}\right) + f\left(\frac{5}{6}\right) \right] \\ &= \frac{1}{3} \left[\left(\frac{1}{6}\right)^2 + \left(\frac{3}{6}\right)^2 + \left(\frac{5}{6}\right)^2 \right] \\ &= \frac{1}{3} \left[\frac{1}{36} + \frac{9}{36} + \frac{25}{36} \right] \\ &= \boxed{\frac{35}{108}} \end{aligned}$$

The estimate Trap(3) is:

$$\begin{aligned} \text{Trap}(3) &= \frac{\Delta x}{2} \left[f(0) + 2f\left(\frac{1}{3}\right) + 2f\left(\frac{2}{3}\right) + f(1) \right] \\ &= \frac{\frac{1}{3}}{2} \left[0^2 + 2\left(\frac{1}{3}\right)^2 + 2\left(\frac{2}{3}\right)^2 + 1^2 \right] \\ &= \frac{1}{6} \left[0 + \frac{2}{9} + \frac{8}{9} + 1 \right] \\ &= \boxed{\frac{19}{54}} \end{aligned}$$

Trap(3) is an overestimate because $f(x)$ is concave up on $[0, 1]$.

**Math 181, Final Exam, Study Guide
Problem 9 Solution**

9. Determine whether the following series converge or not:

$$(a) \sum_{n=1}^{\infty} \frac{n+2}{\sqrt{n^3+n+5}} \quad (b) \sum_{n=1}^{\infty} \frac{(-1)^{n+1}(n^3+1)}{2^n} \quad (c) \sum_{n=1}^{+\infty} \frac{(n^2+1)3^n}{n!} \quad (d) \sum_{n=2}^{+\infty} \frac{1}{n \ln n}$$

Solution:

(a) We note that:

$$0 \leq \frac{n}{\sqrt{n^3+n^3+n^3}} \leq \frac{n+2}{\sqrt{n^3+n+5}}$$

$$0 \leq \frac{1}{\sqrt{3n}} \leq \frac{n+2}{\sqrt{n^3+n+5}}$$

for $n \geq 1$ and that $\sum_{n=1}^{+\infty} \frac{1}{\sqrt{3n}}$ is a divergent p -series with $p = \frac{1}{2} < 1$. Therefore, the series $\sum_{n=1}^{+\infty} \frac{n+2}{\sqrt{n^3+n+5}}$ **diverges** by the Comparison Test.

(b) This series is alternating so we can use the Leibniz Test. The function $\frac{a_n}{2^n} = \frac{n^3+1}{2^n}$ is positive and decreasing for $n \geq 1$. Furthermore,

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{n^3+1}{2^n} = 0$$

Therefore, the series **converges**.

(c) We will use the Ratio Test to show that the series converges.

$$\begin{aligned} \rho &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| \\ &= \lim_{n \rightarrow \infty} \frac{(n+1)^2+1}{(n+1)!} \cdot \frac{n!}{(n^2+1)3^n} \\ &= \lim_{n \rightarrow \infty} \frac{n^2+2n+2}{n^2+1} \cdot \frac{3^{n+1}}{3^n} \cdot \frac{n!}{(n+1)!} \\ &= \lim_{n \rightarrow \infty} \frac{n^2+2n+2}{n^2+1} \cdot 3 \cdot \frac{1}{n+1} \\ &= 3 \lim_{n \rightarrow \infty} \frac{n^2+2n+2}{n^3+n^2+n+1} \\ &= 0 \end{aligned}$$

Since $\rho = 0 < 1$, the series **converges**.

- (d) For this series, we use the Integral Test. This is possible because the function $f(x) = \frac{1}{x \ln x}$ is positive and decreasing for $x \geq 2$. The integral of $f(x)$ on $[2, \infty)$ is:

$$\begin{aligned} \int_2^{+\infty} \frac{dx}{x \ln x} &= \lim_{R \rightarrow +\infty} \int_2^R \frac{dx}{x \ln x} \\ &= \lim_{R \rightarrow +\infty} \left[\ln(\ln x) \right]_2^R \\ &= \lim_{R \rightarrow +\infty} \left[\ln(\ln R) - \ln(\ln 2) \right] \\ &= \infty \end{aligned}$$

Thus, the series **diverges**.

**Math 181, Final Exam, Study Guide
Problem 10 Solution**

10. i) Show that the series

$$\sum_{n=1}^{+\infty} \frac{(-1)^n}{n2^n}$$

converges. ii) Let $S(k)$ be defined by:

$$S(k) = \sum_{n=1}^k \frac{(-1)^n}{n2^n}$$

Then $S(k)$ is the partial sum of the series. Compute k so that $S(k)$ is within .01 of the sum of the series.

Solution: i) The series is alternating so we can use the Leibniz Test. The function $a_n = \frac{1}{n2^n}$ is positive and decreasing for $n \geq 1$. Furthermore,

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{1}{n2^n} = 0$$

Therefore, the series **converges**.

ii) Let S be the exact sum of the series. Then the error $|S - S(k)|$ is bounded as follows:

$$|S - S(k)| \leq a_{k+1}$$

We want the error to be smaller than .01, so we want to choose k to satisfy the following inequality:

$$|S - S(k)| \leq \frac{1}{(k+1)2^{k+1}} < \frac{1}{100}$$

Using a trial and error process, we find that if $k = 3$ then $\frac{1}{(3+1)2^{3+1}} = \frac{1}{64}$ and if $k = 4$ then $\frac{1}{(4+1)2^{4+1}} = \frac{1}{160}$. Thus, we choose $k \geq 4$.

Math 181, Final Exam, Study Guide
Problem 11 Solution

11. Compute the interval of convergence of the following power series:

$$\sum_{n=0}^{+\infty} \frac{(-2)^n (x+4)^n}{n+3} \quad \sum_{n=0}^{+\infty} \frac{3^n (x-1)^{2n}}{n^2}$$

Solution: We use the Ratio Test to find the intervals of convergence. For the first series we have:

$$\begin{aligned} \rho &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{(-2)^{n+1} (x+4)^{n+1}}{(n+1)+3} \cdot \frac{n+3}{2^n (x+4)^n} \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{(-2)^{n+1}}{(-2)^n} \cdot \frac{n+3}{n+4} \cdot \frac{(x+4)^{n+1}}{(x+4)^n} \right| \\ &= \lim_{n \rightarrow \infty} \left| -2 \cdot \frac{n+3}{n+4} \cdot (x+4) \right| \\ &= 2|x+4| \lim_{n \rightarrow \infty} \frac{n+3}{n+4} \\ &= 2|x+4| \cdot (1) \\ &= 2|x+4| \end{aligned}$$

The series converges when $\rho = 2|x+4| < 1$ which gives us:

$$|x+4| < \frac{1}{2} \iff -\frac{1}{2} < x+4 < \frac{1}{2} \iff -\frac{9}{2} < x < -\frac{7}{2}$$

We must now check the endpoints. Plugging $x = -\frac{7}{2}$ into the given power series we get:

$$\sum_{n=0}^{+\infty} \frac{(-2)^n (-\frac{7}{2} + 4)^n}{n+3} = \sum_{n=0}^{+\infty} \frac{(-2)^n (\frac{1}{2})^n}{n+3} = \sum_{n=0}^{+\infty} \frac{(-1)^n}{n+3}$$

which converges by the Leibniz Test.

$$\sum_{n=0}^{+\infty} \frac{(-2)^n (-\frac{9}{2} + 4)^n}{n+3} = \sum_{n=0}^{+\infty} \frac{(-2)^n (-\frac{1}{2})^n}{n+3} = \sum_{n=0}^{+\infty} \frac{1}{n+3}$$

which diverges by a direct comparison with $\sum_{n=1}^{+\infty} \frac{1}{n}$ which is a divergent p -series. Thus, the interval of convergence is:

$$\boxed{-\frac{9}{2} < x \leq -\frac{7}{2}}$$

For the second series we have:

$$\begin{aligned}
 \rho &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| \\
 &= \lim_{n \rightarrow \infty} \left| \frac{3^{n+1}(x-1)^{2(n+1)}}{(n+1)^2} \cdot \frac{n^2}{3^n(x-1)^{2n}} \right| \\
 &= \lim_{n \rightarrow \infty} \left| \frac{3^{n+1}}{3^n} \cdot \frac{n^2}{(n+1)^2} \cdot \frac{(x-1)^{2n+2}}{(x-1)^{2n}} \right| \\
 &= \lim_{n \rightarrow \infty} \left| 3 \cdot \left(\frac{n}{n+1} \right)^2 \cdot (x-1)^2 \right| \\
 &= 3|x-1|^2 \left(\lim_{n \rightarrow \infty} \frac{n}{n+1} \right)^2 \\
 &= 3|x-1|^2 \cdot (1)^2 \\
 &= 3|x-1|^2
 \end{aligned}$$

The series converges when $\rho = 3|x-1|^2 < 1$ which gives us:

$$|x-1|^2 < \frac{1}{3} \iff |x-1| < \frac{1}{\sqrt{3}} \iff -\frac{1}{\sqrt{3}} + 1 < x < \frac{1}{\sqrt{3}} + 1$$

We must now check the endpoints. Plugging $x = \frac{1}{\sqrt{3}} + 1$ into the given power series we get:

$$\sum_{n=1}^{+\infty} \frac{3^n \left(\frac{1}{\sqrt{3}} + 1 - 1 \right)^{2n}}{n^2} = \sum_{n=1}^{+\infty} \frac{3^n \left(\frac{1}{3} \right)^n}{n^2} = \sum_{n=1}^{+\infty} \frac{1}{n^2}$$

which is a convergent p -series ($p = 2 > 1$). Plugging in $x = -\frac{1}{\sqrt{3}} + 1$ we get:

$$\sum_{n=1}^{+\infty} \frac{3^n \left(-\frac{1}{\sqrt{3}} + 1 - 1 \right)^{2n}}{n^2} = \sum_{n=1}^{+\infty} \frac{3^n \left(\frac{1}{3} \right)^n}{n^2} = \sum_{n=1}^{+\infty} \frac{1}{n^2}$$

which, again, is a convergent p -series ($p = 2 > 1$). Thus, the interval of convergence is:

$$\boxed{-\frac{1}{\sqrt{3}} + 1 \leq x \leq \frac{1}{\sqrt{3}} + 1}$$

Math 181, Final Exam, Study Guide
Problem 12 Solution

12. Find the 5th Taylor polynomial of the function $f(x) = \sin(2x)$ centered at $x = 0$.

Solution: We find the 5th Taylor polynomial using the 5th degree Maclaurin polynomial for $\sin(x)$:

$$\sin(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!}$$

and replacing x with $2x$:

$$T_5(x) = (2x) - \frac{(2x)^3}{3!} + \frac{(2x)^5}{5!}$$