

Homogeneous matchbox manifolds

Steven Hurder

University of Illinois at Chicago
www.math.uic.edu/~hurder

Groupoidfest 2009, October 24, 2009

Continua...

Definition: A *continuum* is a compact and connected metrizable space.

Definition: An *indecomposable continuum* is a continuum that is not the union of two proper subcontinua.

Continua...

Definition: A *continuum* is a compact and connected metrizable space.

Definition: An *indecomposable continuum* is a continuum that is not the union of two proper subcontinua.


Examples: The circle \mathbb{S}^1 is decomposable.

Continua...

Definition: A *continuum* is a compact and connected metrizable space.

Definition: An *indecomposable continuum* is a continuum that is not the union of two proper subcontinua.


Examples: The circle \mathbb{S}^1 is decomposable. The Knaster Continuum (or *bucket handle*) is indecomposable.



This is one-half of a Smale Horseshoe. The 2-solenoid over \mathbb{S}^1 is a branched double-covering of it.

Continua...

Indecomposable continua arise naturally as invariant closed sets of dynamical systems; e.g., attractors and minimal sets for diffeomorphisms.



[Picture courtesy Sanjuan, Kennedy, Grebogi & Yorke, "Indecomposable continua in dynamical systems with noise", Chaos 1997]

A Conjecture . . .

Definition: A space X is *homogeneous* if for every $x, y \in X$ there exists a *homeomorphism* $h: X \rightarrow X$ such that $h(x) = y$. Equivalently, X is homogeneous if the group $\text{Homeo}(X)$ acts transitively on X .


Question: [Bing1960] If X is a homogeneous continuum and if every proper subcontinuum of X is an arc, must X then be a circle or a solenoid?

A Conjecture ...

Definition: A space X is *homogeneous* if for every $x, y \in X$ there exists a *homeomorphism* $h: X \rightarrow X$ such that $h(x) = y$. Equivalently, X is homogeneous if the group $\text{Homeo}(X)$ acts transitively on X .

Question: [Bing1960] If X is a homogeneous continuum and if every proper subcontinuum of X is an arc, must X then be a circle or a solenoid?

Theorem: [Hagopian 1977] Let X be a homogeneous continuum such that every proper subcontinuum of X is an arc, then X is an inverse limit over the circle \mathbb{S}^1 .



Matchbox manifolds

Question: Let X be a homogeneous continuum such that every proper subcontinuum of X is an n -dimensional manifold, must X then be an inverse limit of normal coverings of compact manifolds?

Matchbox manifolds

Question: Let X be a homogeneous continuum such that every proper subcontinuum of X is an n -dimensional manifold, must X then be an inverse limit of normal coverings of compact manifolds?

We rephrase the context:


Definition: An n -dimensional *matchbox manifold* is a continuum \mathfrak{M} which is a foliated space with leaf dimension n , and codimension zero.

\mathfrak{M} is a foliated space if it admits a covering $\mathcal{U} = \{\varphi_i \mid 1 \leq i \leq \nu\}$ with foliated coordinate charts $\varphi_i: U_i \rightarrow [-1, 1]^n \times \mathfrak{T}_i$. The compact metric spaces \mathfrak{T}_i are totally disconnected $\iff \mathfrak{M}$ is a matchbox manifold.

The leaves of \mathcal{F} are the path components of \mathfrak{M} .

Smooth matchbox manifolds

Definition: \mathfrak{M} is a *smooth foliated space* if the leafwise transition functions for the foliation charts $\varphi_i: U_i \rightarrow [-1, 1]^n \times \mathfrak{T}_i$ are C^∞ , and vary continuously on the transverse parameter in the leafwise C^∞ -topology.



Automorphisms of matchbox manifolds

A “smooth matchbox manifold” \mathfrak{M} is analogous to a compact manifold, with the transverse dynamics of the foliation \mathcal{F} on the Cantor-like fibers \mathcal{T}_i ; representing fundamental groupoid data. They naturally appear in:

- dynamical systems, as minimal sets & attractors
- geometry, as laminations
- complex dynamics, as universal Riemann surfaces
- algebraic geometry, as models for “stacks”.

Automorphisms of matchbox manifolds

A “smooth matchbox manifold” \mathfrak{M} is analogous to a compact manifold, with the transverse dynamics of the foliation \mathcal{F} on the Cantor-like fibers \mathcal{T}_i ; representing fundamental groupoid data. They naturally appear in:

- dynamical systems, as minimal sets & attractors
- geometry, as laminations
- complex dynamics, as universal Riemann surfaces
- algebraic geometry, as models for “stacks”.

Bing Question: For which \mathfrak{M} is the group $\text{Homeo}(\mathfrak{M})$ transitive?

Automorphisms of matchbox manifolds

A “smooth matchbox manifold” \mathfrak{M} is analogous to a compact manifold, with the transverse dynamics of the foliation \mathcal{F} on the Cantor-like fibers \mathcal{T}_i ; representing fundamental groupoid data. They naturally appear in:

- dynamical systems, as minimal sets & attractors
- geometry, as laminations
- complex dynamics, as universal Riemann surfaces
- algebraic geometry, as models for “stacks”.

Bing Question: For which \mathfrak{M} is the group $\text{Homeo}(\mathfrak{M})$ transitive?

Klein Question: Do the Riemannian symmetries of \mathfrak{M} characterize it?

Automorphisms of matchbox manifolds

A “smooth matchbox manifold” \mathfrak{M} is analogous to a compact manifold, with the transverse dynamics of the foliation \mathcal{F} on the Cantor-like fibers \mathcal{T}_i ; representing fundamental groupoid data. They naturally appear in:

- dynamical systems, as minimal sets & attractors
- geometry, as laminations
- complex dynamics, as universal Riemann surfaces
- algebraic geometry, as models for “stacks”.

Bing Question: For which \mathfrak{M} is the group $\text{Homeo}(\mathfrak{M})$ transitive?

Klein Question: Do the Riemannian symmetries of \mathfrak{M} characterize it?

Zimmer Question: What countable groups Λ act effectively on \mathfrak{M} ?

Automorphisms of matchbox manifolds

A “smooth matchbox manifold” \mathfrak{M} is analogous to a compact manifold, with the transverse dynamics of the foliation \mathcal{F} on the Cantor-like fibers \mathcal{T}_i ; representing fundamental groupoid data. They naturally appear in:

- dynamical systems, as minimal sets & attractors
- geometry, as laminations
- complex dynamics, as universal Riemann surfaces
- algebraic geometry, as models for “stacks”.

Bing Question: For which \mathfrak{M} is the group $\text{Homeo}(\mathfrak{M})$ transitive?

Klein Question: Do the Riemannian symmetries of \mathfrak{M} characterize it?

Zimmer Question: What countable groups Λ act effectively on \mathfrak{M} ?

Haefliger Question: What are the topological invariants associated to matchbox manifolds, and do they characterize them in some fashion?

A solution to the Bing Question

Theorem [Clark & Hurder 2009] Let \mathfrak{M} be an orientable homogeneous smooth matchbox manifold. Then \mathfrak{M} is homeomorphic to a McCord (or normal) solenoid. In particular, \mathfrak{M} is minimal, so every leaf is dense.

A solution to the Bing Question

Theorem [Clark & Hurder 2009] Let \mathfrak{M} be an orientable homogeneous smooth matchbox manifold. Then \mathfrak{M} is homeomorphic to a McCord (or normal) solenoid. In particular, \mathfrak{M} is minimal, so every leaf is dense.

When the dimension of \mathfrak{M} is $n = 1$ (that is, \mathcal{F} is defined by a flow) then this recovers the result of Hagopian, but the proof is much closer in spirit to the later proof of this case by [Aarts, Hagopian and Oversteegen 1991].

The case where \mathfrak{M} is given as a fibration over \mathbb{T}^n with totally disconnected fibers was proven in [Clark, 2002].

The key to the proof in the general case is the extensive use of *pseudogroups* and *groupoids* – in place of Lie group actions.

Two applications

Here are two consequences of the Main Theorem:

Corollary: Let \mathfrak{M} be an orientable homogeneous n -dimensional smooth matchbox manifold, which is embedded in a closed $(n + 1)$ -dimensional manifold. Then \mathfrak{M} is itself a manifold.

For \mathfrak{M} a homogeneous continuum with a non-singular flow, this was a question/conjecture of Bing, solved by [Thomas 1971]. Non-embedding for solenoids of dimension $n \geq 2$ was solved by [Clark & Fokkink, 2002]. Proofs use shape theory and Alexander-Spanier duality for cohomology.

Two applications

Here are two consequences of the Main Theorem:

Corollary: Let \mathfrak{M} be an orientable homogeneous n -dimensional smooth matchbox manifold, which is embedded in a closed $(n + 1)$ -dimensional manifold. Then \mathfrak{M} is itself a manifold.

For \mathfrak{M} a homogeneous continuum with a non-singular flow, this was a question/conjecture of Bing, solved by [Thomas 1971]. Non-embedding for solenoids of dimension $n \geq 2$ was solved by [Clark & Fokkink, 2002]. Proofs use shape theory and Alexander-Spanier duality for cohomology.

Corollary: Let \mathfrak{M} be the tiling space associated to a tiling \mathcal{P} of \mathbb{R}^n . If \mathfrak{M} is homogeneous, then the tiling is periodic.

Generalized solenoids

Let M_ℓ be compact, orientable manifolds of dimension $n \geq 1$ for $\ell \geq 0$, with orientation-preserving covering maps

$$\xrightarrow{p_{\ell+1}} M_\ell \xrightarrow{p_\ell} M_{\ell-1} \xrightarrow{p_{\ell-1}} \cdots \xrightarrow{p_2} M_1 \xrightarrow{p_1} M_0$$

Generalized solenoids

Let M_ℓ be compact, orientable manifolds of dimension $n \geq 1$ for $\ell \geq 0$, with orientation-preserving covering maps

$$\xrightarrow{p_{\ell+1}} M_\ell \xrightarrow{p_\ell} M_{\ell-1} \xrightarrow{p_{\ell-1}} \cdots \xrightarrow{p_2} M_1 \xrightarrow{p_1} M_0$$

The p_ℓ are called the *bonding maps* for the solenoid

$$\mathcal{S} = \varprojlim \{p_\ell: M_\ell \rightarrow M_{\ell-1}\} \subset \prod_{\ell=0}^{\infty} M_\ell$$

Generalized solenoids

Let M_ℓ be compact, orientable manifolds of dimension $n \geq 1$ for $\ell \geq 0$, with orientation-preserving covering maps

$$\xrightarrow{p_{\ell+1}} M_\ell \xrightarrow{p_\ell} M_{\ell-1} \xrightarrow{p_{\ell-1}} \cdots \xrightarrow{p_2} M_1 \xrightarrow{p_1} M_0$$

The p_ℓ are called the *bonding maps* for the solenoid

$$\mathcal{S} = \varprojlim \{p_\ell: M_\ell \rightarrow M_{\ell-1}\} \subset \prod_{\ell=0}^{\infty} M_\ell$$

Choose basepoints $x_\ell \in M_\ell$ with $p_\ell(x_\ell) = x_{\ell-1}$. Set $G_\ell = \pi_1(M_\ell, x_\ell)$.

Then we have a descending chain of groups and injective maps

$$\xrightarrow{p_{\ell+1}} G_\ell \xrightarrow{p_\ell} G_{\ell-1} \xrightarrow{p_{\ell-1}} \cdots \xrightarrow{p_2} G_1 \xrightarrow{p_1} G_0$$

Set $q_\ell = p_\ell \circ \cdots \circ p_1: M_\ell \longrightarrow M_0$.

McCord solenoids

Definition: S is a *McCord solenoid* for some fixed $\ell_0 \geq 0$, for all $\ell \geq \ell_0$ the image H_ℓ of G_ℓ in $H_{\ell_0} \equiv G_{\ell_0}$ is a normal subgroup.

Theorem [McCord 1965] A McCord solenoid S is an orientable homogeneous smooth matchbox manifold.

McCord solenoids

Definition: \mathcal{S} is a *McCord solenoid* for some fixed $\ell_0 \geq 0$, for all $\ell \geq \ell_0$ the image H_ℓ of G_ℓ in $H_{\ell_0} \equiv G_{\ell_0}$ is a normal subgroup.

Theorem [McCord 1965] A McCord solenoid \mathcal{S} is an orientable homogeneous smooth matchbox manifold.

Remark: $\pi_1(M_0, x_0)$ nilpotent implies that \mathcal{S} is a McCord solenoid.

McCord solenoids

Definition: \mathcal{S} is a *McCord solenoid* for some fixed $\ell_0 \geq 0$, for all $\ell \geq \ell_0$ the image H_ℓ of G_ℓ in $H_{\ell_0} \equiv G_{\ell_0}$ is a normal subgroup.

Theorem [McCord 1965] A McCord solenoid \mathcal{S} is an orientable homogeneous smooth matchbox manifold.

Remark: $\pi_1(M_0, x_0)$ nilpotent implies that \mathcal{S} is a McCord solenoid.

Caution: There are constructions of inverse limits \mathcal{S} as above where the bonding maps are not normal coverings, and the McCord condition does not hold, but \mathcal{S} is homogeneous [Fokkink & Oversteegen 2002].

Our technique of proof of the main theorem for such examples presents the inverse limit space \mathcal{S} as homeomorphic to a normal tower of coverings.

Effros Theorem

Let X be a separable and metrizable topological space. Let G be a topological group with identity e .

For $U \subseteq G$ and $x \in X$, let $Ux = \{gx \mid g \in U\}$.

Definition: An action of G on X is *transitive* if $Gx = X$ for all $x \in X$.

Effros Theorem

Let X be a separable and metrizable topological space. Let G be a topological group with identity e .

For $U \subseteq G$ and $x \in X$, let $Ux = \{gx \mid g \in U\}$.

Definition: An action of G on X is *transitive* if $Gx = X$ for all $x \in X$.

Definition: An action of G on X is *micro-transitive* if for every $x \in X$ and every neighborhood U of e , Ux is a neighborhood of x .

Effros Theorem

Let X be a separable and metrizable topological space. Let G be a topological group with identity e .

For $U \subseteq G$ and $x \in X$, let $Ux = \{gx \mid g \in U\}$.

Definition: An action of G on X is *transitive* if $Gx = X$ for all $x \in X$.

Definition: An action of G on X is *micro-transitive* if for every $x \in X$ and every neighborhood U of e , Ux is a neighborhood of x .

Theorem [Effros 1965] Suppose that a completely metrizable group G acts *transitively* on a second category space X , then it acts *micro-transitively* on X .

Effros Theorem

Let X be a separable and metrizable topological space. Let G be a topological group with identity e .

For $U \subseteq G$ and $x \in X$, let $Ux = \{gx \mid g \in U\}$.

Definition: An action of G on X is *transitive* if $Gx = X$ for all $x \in X$.

Definition: An action of G on X is *micro-transitive* if for every $x \in X$ and every neighborhood U of e , Ux is a neighborhood of x .

Theorem [Effros 1965] Suppose that a completely metrizable group G acts *transitively* on a second category space X , then it acts *micro-transitively* on X .

Alternate proofs of have been given by [Ancel 1987] and [van Mill 2004]. Remarkably, Van Mill shows that Effros Theorem is equivalent to the *Open Mapping Principle* of Functional Analysis. This appeared in the American Mathematical Monthly, pages 801–806, 2004.

Interpretation for compact metric spaces

The metric on the group $\text{Homeo}(X)$ for (X, d_X) a separable, locally compact, metric space is given by

$$d_H(f, g) := \sup \{d_X(f(x), g(x)) \mid x \in X\} \\ + \sup \{d_X(f^{-1}(x), g^{-1}(x)) \mid x \in X\}$$

Interpretation for compact metric spaces

The metric on the group $\text{Homeo}(X)$ for (X, d_X) a separable, locally compact, metric space is given by

$$d_H(f, g) := \sup \{d_X(f(x), g(x)) \mid x \in X\} \\ + \sup \{d_X(f^{-1}(x), g^{-1}(x)) \mid x \in X\}$$

Corollary: Let X be a homogeneous compact metric space. Then for any given $\epsilon > 0$ there is a corresponding $\delta > 0$ so that if $d_X(x, y) < \delta$, there is a homeomorphism $h: X \rightarrow X$ with $d_H(h, id_X) < \epsilon$ and $h(x) = y$.

In particular, for a homogeneous foliated space \mathfrak{M} this conclusion holds.

This observation was used by [Aarts, Hagopian, & Oversteegen 1991] and [Clark 2002] in their study of matchbox manifolds.

Holonomy groupoids

Let $\varphi_i: U_i \rightarrow [-1, 1]^n \times \mathfrak{F}_i$ for $1 \leq i \leq \nu$ be the covering of \mathfrak{M} by foliation charts. For $U_i \cap U_j \neq \emptyset$ we obtain the holonomy transformation

$$h_{ji}: D(h_{ji}) \subset \mathfrak{F}_i \longrightarrow R(h_{ji}) \subset \mathfrak{F}_j$$

These transformations generate the holonomy pseudogroup $\mathcal{G}_{\mathcal{F}}$ of \mathfrak{M} , modeled on the transverse metric space $\mathfrak{F} = \mathfrak{F}_1 \cup \dots \cup \mathfrak{F}_{\nu}$

Typical element of $\mathcal{G}_{\mathcal{F}}$ is a composition, for $\mathcal{I} = (i_0, i_1, \dots, i_k)$ where $U_{i_\ell} \cap U_{i_{\ell-1}} \neq \emptyset$ for $1 \leq \ell \leq k$,

$$h_{\mathcal{I}} = h_{i_k i_{k-1}} \circ \dots \circ h_{i_1 i_0}: D(h_{\mathcal{I}}) \subset \mathfrak{F}_{i_0} \longrightarrow R(h_{\mathcal{I}}) \subset \mathfrak{F}_{i_k}$$

$x \in \mathfrak{F}$ is a *point of holonomy* for $\mathcal{G}_{\mathcal{F}}$ if there exists some $h_{\mathcal{I}} \in \mathcal{G}_{\mathcal{F}}$ with $x \in D(h_{\mathcal{I}})$ such that $h_{\mathcal{I}}(x) = x$ and the germ of $h_{\mathcal{I}}$ at x is non-trivial.

We say \mathcal{F} is *without holonomy* if there are no points of holonomy.

Equicontinuous matchbox manifolds

Definition: \mathfrak{M} is an *equicontinuous matchbox manifold* if it admits some covering by foliation charts as above, such that for all $\epsilon > 0$, there exists $\delta > 0$ so that for all $h_{\mathcal{I}} \in \mathcal{G}_{\mathcal{F}}$ we have

$$x, y \in D(h_{\mathcal{I}}) \text{ with } d_{\mathfrak{X}}(x, y) < \delta \implies d_{\mathfrak{X}}(h_{\mathcal{I}}(x), h_{\mathcal{I}}(y)) < \epsilon$$

Equicontinuous matchbox manifolds

Definition: \mathfrak{M} is an *equicontinuous matchbox manifold* if it admits some covering by foliation charts as above, such that for all $\epsilon > 0$, there exists $\delta > 0$ so that for all $h_{\mathcal{I}} \in \mathcal{G}_{\mathcal{F}}$ we have

$$x, y \in D(h_{\mathcal{I}}) \text{ with } d_{\mathfrak{X}}(x, y) < \delta \implies d_{\mathfrak{X}}(h_{\mathcal{I}}(x), h_{\mathcal{I}}(y)) < \epsilon$$

Theorem: A homogeneous matchbox manifold \mathfrak{M} is equicontinuous without holonomy.

Equicontinuous matchbox manifolds

Definition: \mathfrak{M} is an *equicontinuous matchbox manifold* if it admits some covering by foliation charts as above, such that for all $\epsilon > 0$, there exists $\delta > 0$ so that for all $h_I \in \mathcal{G}_{\mathcal{F}}$ we have

$$x, y \in D(h_I) \text{ with } d_{\mathfrak{X}}(x, y) < \delta \implies d_{\mathfrak{X}}(h_I(x), h_I(y)) < \epsilon$$

Theorem: A homogeneous matchbox manifold \mathfrak{M} is equicontinuous without holonomy.

The proof relies on one basic observation and *extensive* technical analysis.

Lemma: Let $h: \mathfrak{M} \rightarrow \mathfrak{M}$ be a homeomorphism. Then h maps the leaves of \mathcal{F} to leaves of \mathcal{F} . That is, every $h \in \text{Homeo}(\mathfrak{M})$ is foliation-preserving.

Proof: The leaves of \mathcal{F} are the path components of \mathfrak{M} .

Equicontinuous matchbox manifolds

Definition: \mathfrak{M} is an *equicontinuous matchbox manifold* if it admits some covering by foliation charts as above, such that for all $\epsilon > 0$, there exists $\delta > 0$ so that for all $h_I \in \mathcal{G}_{\mathcal{F}}$ we have

$$x, y \in D(h_I) \text{ with } d_{\mathfrak{X}}(x, y) < \delta \implies d_{\mathfrak{X}}(h_I(x), h_I(y)) < \epsilon$$

Theorem: A homogeneous matchbox manifold \mathfrak{M} is equicontinuous without holonomy.

The proof relies on one basic observation and *extensive* technical analysis.

Lemma: Let $h: \mathfrak{M} \rightarrow \mathfrak{M}$ be a homeomorphism. Then h maps the leaves of \mathcal{F} to leaves of \mathcal{F} . That is, every $h \in \text{Homeo}(\mathfrak{M})$ is foliation-preserving.

Proof: The leaves of \mathcal{F} are the path components of \mathfrak{M} .

Theorem: An equicontinuous matchbox manifold \mathfrak{M} is minimal.

Three Structure Theorems

We can now state the three main structure theorems.

Theorem 1: Let \mathfrak{M} be an equicontinuous matchbox manifold without holonomy. Then \mathfrak{M} is homeomorphic to a solenoid

$$\mathcal{S} = \varprojlim \{p_\ell: M_\ell \rightarrow M_{\ell-1}\}$$

Three Structure Theorems

We can now state the three main structure theorems.

Theorem 1: Let \mathfrak{M} be an equicontinuous matchbox manifold without holonomy. Then \mathfrak{M} is homeomorphic to a solenoid

$$\mathcal{S} = \varprojlim \{p_\ell: M_\ell \rightarrow M_{\ell-1}\}$$

Theorem 2: Let \mathfrak{M} be a homogeneous matchbox manifold. Then the bonding maps above can be chosen so that $q_\ell: M_\ell \rightarrow M_0$ is a normal covering for all $\ell \geq 0$. That is, \mathcal{S} is McCord.

Three Structure Theorems

We can now state the three main structure theorems.

Theorem 1: Let \mathfrak{M} be an equicontinuous matchbox manifold without holonomy. Then \mathfrak{M} is homeomorphic to a solenoid

$$\mathcal{S} = \varprojlim \{p_\ell: M_\ell \rightarrow M_{\ell-1}\}$$

Theorem 2: Let \mathfrak{M} be a homogeneous matchbox manifold. Then the bonding maps above can be chosen so that $q_\ell: M_\ell \rightarrow M_0$ is a normal covering for all $\ell \geq 0$. That is, \mathcal{S} is McCord.

Theorem 3: Let \mathfrak{M} be a homogeneous matchbox manifold. Then there exists a clopen subset $V \subset \mathfrak{T}$ such that the restricted groupoid $\mathcal{H}(\mathcal{F}, V) \equiv \mathcal{G}_{\mathcal{F}}|_V$ is a group, and \mathfrak{M} is homeomorphic to the suspension of the action of $\mathcal{H}(\mathcal{F}, V)$ on V . Thus, the fibers of the map $q_\infty: \mathfrak{M} \rightarrow M_0$ are homeomorphic to a profinite completion of $\mathcal{H}(\mathcal{F}, V)$.

Coding & Quasi-Tiling

Let \mathfrak{M} be an equicontinuous matchbox manifold without holonomy.

Fix basepoint $w_0 \in \text{int}(\mathfrak{T}_1)$ with corresponding leaf $L_0 \subset \mathfrak{M}$.

The equivalence relation on \mathfrak{T} induced by \mathcal{F} is denoted Γ , and we have the following subsets:

$$\Gamma_W = \{(w, w') \mid w \in W, w' \in \mathcal{O}(w)\}$$

$$\Gamma_W^W = \{(w, w') \mid w \in W, w' \in \mathcal{O}(w) \cap W\}$$

$$\Gamma_0 = \{w' \in W \mid (w_0, w') \in \Gamma_W^W\} = \mathcal{O}(w_0) \cap W$$

Note that Γ_W^W is a groupoid, with object space W . The assumption that \mathcal{F} is without holonomy implies Γ_W^W is equivalent to the groupoid of germs of local holonomy maps induced from the restriction of $\mathcal{G}_{\mathcal{F}}$ to W .

Equicontinuity & uniform domains

Proposition: Let \mathfrak{M} be an equicontinuous matchbox manifold without holonomy. Given $\epsilon_* > 0$, then there exists $\delta_* > 0$ such that:

- for all $(w, w') \in \Gamma_W^W$ the corresponding holonomy map $h_{w,w'}$ satisfies $D_{\mathfrak{I}}(w, \delta_*) \subset D(h_{w,w'})$
- $d_{\mathfrak{I}}(h_{w,w'}(z), h_{w,w'}(z')) < \epsilon_*$ for all $z, z' \in D_{\mathfrak{I}}(w, \delta_*)$.

Equicontinuity & uniform domains

Proposition: Let \mathfrak{M} be an equicontinuous matchbox manifold without holonomy. Given $\epsilon_* > 0$, then there exists $\delta_* > 0$ such that:

- for all $(w, w') \in \Gamma_W^W$ the corresponding holonomy map $h_{w,w'}$ satisfies $D_{\mathfrak{T}}(w, \delta_*) \subset D(h_{w,w'})$
- $d_{\mathfrak{T}}(h_{w,w'}(z), h_{w,w'}(z')) < \epsilon_*$ for all $z, z' \in D_{\mathfrak{T}}(w, \delta_*)$.

Let $W \subset \mathfrak{T}_1$ be a clopen subset with $w_0 \in W$. Decompose W into clopen subsets of diameter $\epsilon_\ell > 0$,

$$W = W_1^\ell \cup \dots \cup W_{\beta_\ell}^\ell$$

Set $\eta_\ell = \min \left\{ d_{\mathfrak{T}}(W_i^\ell, W_j^\ell) \mid 1 \leq i \neq j \leq \beta_\ell \right\} > 0$ and let $\delta_\ell > 0$ be the constant of equicontinuity as above.

The orbit coding function

- The code space $\mathcal{C}_\ell = \{1, \dots, \beta_\ell\}$
- For $w \in W$, the \mathcal{C}_w^ℓ -code of $u \in W$ is the function $C_{w,u}^\ell: \Gamma_w \rightarrow \mathcal{C}_\ell$ defined as: for $w' \in \Gamma_w$ set $C_{w,u}^\ell(w') = i$ if $h_{w,w'}(u) \in W_i^\ell$.
- Define $V^\ell = \left\{ u \in W_1^\ell \mid C_{w_0,u}^\ell(w') = C_{w_0,w_0}^\ell(w') \text{ for all } w' \in \Gamma_0 \right\}$

The orbit coding function

- The code space $\mathcal{C}_\ell = \{1, \dots, \beta_\ell\}$
- For $w \in W$, the \mathcal{C}_w^ℓ -code of $u \in W$ is the function $C_{w,u}^\ell: \Gamma_w \rightarrow \mathcal{C}_\ell$ defined as: for $w' \in \Gamma_w$ set $C_{w,u}^\ell(w') = i$ if $h_{w,w'}(u) \in W_i^\ell$.
- Define $V^\ell = \left\{ u \in W_1^\ell \mid C_{w_0,u}^\ell(w') = C_{w_0,w_0}^\ell(w') \text{ for all } w' \in \Gamma_0 \right\}$

Lemma: If $u, v \in W$ with $d_{\mathfrak{X}}(u, v) < \delta_\ell$ then $C_{w,u}^\ell(w') = C_{w,v}^\ell(w')$ for all $w' \in \Gamma_w$. Hence, the function $C_w^\ell(u) = C_{w,u}^\ell$ is locally constant in u .

Thus, V^ℓ is open, and the translates of this set define a Γ_0 -invariant clopen decomposition of W .

The coding decomposition

The Thomas tube $\tilde{\mathfrak{M}}_\ell$ for \mathfrak{M} is the “saturation” of V^ℓ by \mathcal{F} .

The saturation is necessarily all of \mathfrak{M} . But the tube structure comes with a vertical fibration, which allows for collapsing the tube in foliation charts. This is the basis of the main technical result:

The coding decomposition

The Thomas tube $\tilde{\mathfrak{M}}_\ell$ for \mathfrak{M} is the “saturation” of V^ℓ by \mathcal{F} .

The saturation is necessarily all of \mathfrak{M} . But the tube structure comes with a vertical fibration, which allows for collapsing the tube in foliation charts. This is the basis of the main technical result:

Theorem: For $\text{diam}(V^\ell)$ sufficiently small, there is a quotient map $\Pi_\ell: \tilde{\mathfrak{M}}_\ell \rightarrow M_\ell$ whose fibers are the transversal sections isotopic to V^ℓ , and whose base is a compact manifold. This yields compatible maps $\Pi_\ell: \mathfrak{M} \rightarrow M_\ell$ which induce the solenoid structure on \mathfrak{M} .

Furthermore, if \mathfrak{M} is homogeneous, then $\text{Homeo}(\mathfrak{M})$ acts transitively on the fibers of the tower induced by the maps $\Pi_\ell: \mathfrak{M} \rightarrow M_\ell$, hence the tower is normal.

Leeuwenbrug Conjecture

Conjecture: Let \mathfrak{M} be an equicontinuous matchbox manifold, and $V \subset \mathfrak{T}$ a clopen set. Then \mathfrak{M} is characterized up to homeomorphism by the restricted groupoid $\mathcal{H}(\mathcal{F}, V) \equiv \mathcal{G}_{\mathcal{F}}|V$ and any partial quotient M_{ℓ} .

That is, for matchbox manifolds, Kakutani equivalence implies homeomorphism (modulo some obvious additional conditions.)

Leeuwenbrug Conjecture

Conjecture: Let \mathfrak{M} be an equicontinuous matchbox manifold, and $V \subset \mathfrak{T}$ a clopen set. Then \mathfrak{M} is characterized up to homeomorphism by the restricted groupoid $\mathcal{H}(\mathcal{F}, V) \equiv \mathcal{G}_{\mathcal{F}}|V$ and any partial quotient M_{ℓ} .

That is, for matchbox manifolds, Kakutani equivalence implies homeomorphism (modulo some obvious additional conditions.)

This is known for flows [Dye 1957, Fokkink 1991].

Leeuwenbrug Conjecture

Conjecture: Let \mathfrak{M} be an equicontinuous matchbox manifold, and $V \subset \mathfrak{T}$ a clopen set. Then \mathfrak{M} is characterized up to homeomorphism by the restricted groupoid $\mathcal{H}(\mathcal{F}, V) \equiv \mathcal{G}_{\mathcal{F}}|V$ and any partial quotient M_{ℓ} .

That is, for matchbox manifolds, Kakutani equivalence implies homeomorphism (modulo some obvious additional conditions.)

This is known for flows [Dye 1957, Fokkink 1991].



Happy Birthday, Jean!



Jean Renault - Boulder 1999