

# Dynamics and Cohomology of Foliations


Steven Hurder

University of Illinois at Chicago  
[www.math.uic.edu/~hurder](http://www.math.uic.edu/~hurder)

VIII International Colloquium on Differential Geometry  
Santiago de Compostela, 7-11 July 2008

## Definition of foliation

A foliation  $\mathcal{F}$  of dimension  $p$  on a manifold  $M^m$  is a decomposition into “uniform layers” – the leaves – which are immersed submanifolds of codimension  $q$ : there is an open covering of  $M$  by coordinate charts so that the leaves are mapped into linear planes of dimension  $p$ , and the transition function preserves these planes.



# Fundamental problems

**Problem:** “Classify” the foliations on a given manifold  $M$

# Fundamental problems

**Problem:** “Classify” the foliations on a given manifold  $M$

Two classification schemes have been developed since 1970:  
using either “homotopy” or “dynamics”.

**Question:** How are the cohomology invariants of a foliation related to its dynamical behavior?

# Integrable homotopy equivalence

Let  $q$  denote the codimension of the foliation  $\mathcal{F}$ .

$q = m - p$  where  $p$  is the leaf dimension,  $m = \dim M$

Assume throughout that  $\mathcal{F}$  is transversally  $C^r$  for  $r \geq 2$ .

# Integrable homotopy equivalence

Let  $q$  denote the codimension of the foliation  $\mathcal{F}$ .

$q = m - p$  where  $p$  is the leaf dimension,  $m = \dim M$

Assume throughout that  $\mathcal{F}$  is transversally  $C^r$  for  $r \geq 2$ .

**Definition:** Two foliations  $\mathcal{F}_0$  and  $\mathcal{F}_1$  of codimension  $q$  on  $M$  are *integrably homotopic* if there exists a foliation  $\mathcal{F}$  of codimension  $q$  on  $M \times \mathbb{R}$  which is transverse to the slices  $M \times \{t\} \subset M \times \mathbb{R}$  for  $t = 0, 1$ , such that the restrictions  $\mathcal{F}|_{M \times \{t\}} = \mathcal{F}_t$  for  $t = 0, 1$ .

Integrable homotopy is a fairly weak notion of equivalence. For example, if  $M$  is an open contractible manifold then any two foliations  $\mathcal{F}_0$  and  $\mathcal{F}_1$  on  $M$  are integrably homotopic.

## Classifying spaces:

$B\Gamma_q$  denotes the “classifying space” of codimension- $q$  foliations introduced by André Haefliger in 1970.

There is a natural map  $\nu: B\Gamma_q \rightarrow BO_q$ .

## Classifying spaces:

$B\Gamma_q$  denotes the “classifying space” of codimension- $q$  foliations introduced by André Haefliger in 1970.

There is a natural map  $\nu: B\Gamma_q \rightarrow BO_q$ .

**Theorem:** (Haefliger [1970]) Each foliation  $\mathcal{F}$  on  $M$  of codimension  $q$  determines a well-defined map  $h_{\mathcal{F}}: M \rightarrow B\Gamma_q$  whose homotopy class is uniquely defined by  $\mathcal{F}$ , and depends only upon the integrable homotopy class of  $\mathcal{F}$ . The composition  $\nu \circ h_{\mathcal{F}}: M \rightarrow BO_q$  classifies the normal bundle  $Q \rightarrow M$  of  $\mathcal{F}$ .



## Classifying spaces:

$B\Gamma_q$  denotes the “classifying space” of codimension- $q$  foliations introduced by André Haefliger in 1970.

There is a natural map  $\nu: B\Gamma_q \rightarrow BO_q$ .

**Theorem:** (Haefliger [1970]) Each foliation  $\mathcal{F}$  on  $M$  of codimension  $q$  determines a well-defined map  $h_{\mathcal{F}}: M \rightarrow B\Gamma_q$  whose homotopy class is uniquely defined by  $\mathcal{F}$ , and depends only upon the integrable homotopy class of  $\mathcal{F}$ . The composition  $\nu \circ h_{\mathcal{F}}: M \rightarrow BO_q$  classifies the normal bundle  $Q \rightarrow M$  of  $\mathcal{F}$ .

**Theorem:** (Thurston [1975]) Let  $M$  be a closed manifold. A map  $h: M \rightarrow B\Gamma_q \times BO_p$  for which the composition

$$(\nu \times Id) \circ h: M \rightarrow BO_q \times BO_p \rightarrow BO_m$$

classifies the tangent bundle  $TM$ , determines an integrable homotopy class of a codimension- $q$  foliation  $\mathcal{F}_h$  on  $M$ .

## Primary characteristic classes

The Pontrjagin classes of the normal bundle  $Q \rightarrow M$  factor through the map:

$$\begin{array}{ccc} & & B\Gamma_q \\ & \nearrow h_{\mathcal{F}} & \downarrow \nu \\ M & \xrightarrow{h_Q} & BO_q \end{array}$$

## Primary characteristic classes

The Pontrjagin classes of the normal bundle  $Q \rightarrow M$  factor through the map:

$$\begin{array}{ccc} & & B\Gamma_q \\ & \nearrow h_{\mathcal{F}} & \downarrow \nu \\ M & \xrightarrow{h_Q} & BO_q \end{array}$$

**Theorem:** (Bott [1970])

$h_Q^*: H^\ell(BO_q; \mathbb{R}) \rightarrow H^\ell(M; \mathbb{R})$  is trivial for  $\ell > 2q$ .

## Primary characteristic classes

The Pontrjagin classes of the normal bundle  $Q \rightarrow M$  factor through the map:

$$\begin{array}{ccc} & & B\Gamma_q \\ & \nearrow h_{\mathcal{F}} & \downarrow \nu \\ M & \xrightarrow{h_Q} & BO_q \end{array}$$

**Theorem:** (Bott [1970])

$$h_Q^*: H^\ell(BO_q; \mathbb{R}) \rightarrow H^\ell(M; \mathbb{R}) \text{ is trivial for } \ell > 2q.$$

**Theorem:** (Bott-Heitsch [1972])

$$h_Q^*: H^\ell(BO_q; \mathbb{Z}) \rightarrow H^\ell(M; \mathbb{Z}) \text{ is injective for all } \ell.$$

## Secondary classes

**Theorem:** (Godbillon-Vey [1971]) For  $q \geq 1$ , the Godbillon-Vey class  $GV(\mathcal{F}) = \Delta(h_1 c_1^q) \in H^{2q+1}(M; \mathbb{R})$  is an integrable homotopy invariant.

## Secondary classes

**Theorem:** (Godbillon-Vey [1971]) For  $q \geq 1$ , the Godbillon-Vey class  $GV(\mathcal{F}) = \Delta(h_1 c_1^q) \in H^{2q+1}(M; \mathbb{R})$  is an integrable homotopy invariant.

$$WO_q \cong \Lambda(h_1, h_3, \dots, h_{q/2}) \otimes \mathbb{R}[c_1, c_2, \dots, c_q], \quad d_W h_i = c_i, \quad d_W c_i = 0$$

**Theorem:** (Bott-Haefliger, Gelfand-Fuks, Kamber-Tondeur [1972]) For  $q \geq 1$ , there is a non-trivial space of secondary invariants  $H^*(WO_q)$  and functorial characteristic map whose image contains the Godbillon-Vey class

$$\begin{array}{ccc} & & H^*(B\Gamma_q; \mathbb{R}) \\ & \nearrow \tilde{\Delta} & \downarrow h_{\mathcal{F}}^* \\ H^*(WO_q) & \xrightarrow{\Delta_{\mathcal{F}}} & H^*(M; \mathbb{R}) \end{array}$$

The study of the images of the maps  $\Delta_{\mathcal{F}}$  has been the principle source of information about the (non-trivial) homotopy type of  $B\Gamma_q$ .

# Homotopy chaos

**Theorem:** (Bott–Heitsch [1972])  $B\Gamma_q^r$  does not have finite topological type for  $q \geq 2$ .

# Homotopy chaos

**Theorem:** (Bott–Heitsch [1972])  $B\Gamma_q^r$  does not have finite topological type for  $q \geq 2$ .

**Theorem:** (Thurston [1972])  $\pi_3(B\Gamma_1^r)$  surjects onto  $\mathbb{R}$ .



# Homotopy chaos

**Theorem:** (Bott–Heitsch [1972])  $B\Gamma_q^r$  does not have finite topological type for  $q \geq 2$ .

**Theorem:** (Thurston [1972])  $\pi_3(B\Gamma_1^r)$  surjects onto  $\mathbb{R}$ .

**Theorem:** (Heitsch [1978]) There are continuous families of foliations with non-trivial variations of their secondary classes for  $q \geq 3$ .

# Homotopy chaos

**Theorem:** (Bott–Heitsch [1972])  $B\Gamma_q^r$  does not have finite topological type for  $q \geq 2$ .

**Theorem:** (Thurston [1972])  $\pi_3(B\Gamma_1^r)$  surjects onto  $\mathbb{R}$ .

**Theorem:** (Heitsch [1978]) There are continuous families of foliations with non-trivial variations of their secondary classes for  $q \geq 3$ .

**Theorem:** (Rasmussen [1980]) There are continuous families of foliations with non-trivial variations of their secondary classes for  $q = 2$ .

# Homotopy chaos

**Theorem:** (Bott–Heitsch [1972])  $B\Gamma_q^r$  does not have finite topological type for  $q \geq 2$ .

**Theorem:** (Thurston [1972])  $\pi_3(B\Gamma_1^r)$  surjects onto  $\mathbb{R}$ .

**Theorem:** (Heitsch [1978]) There are continuous families of foliations with non-trivial variations of their secondary classes for  $q \geq 3$ .

**Theorem:** (Rasmussen [1980]) There are continuous families of foliations with non-trivial variations of their secondary classes for  $q = 2$ .

**Corollary:**  $B\Gamma_q^r$  has uncountable topological type for all  $q \geq 1$ .

# Homotopy chaos

**Theorem:** (Bott–Heitsch [1972])  $B\Gamma_q^r$  does not have finite topological type for  $q \geq 2$ .

**Theorem:** (Thurston [1972])  $\pi_3(B\Gamma_1^r)$  surjects onto  $\mathbb{R}$ .

**Theorem:** (Heitsch [1978]) There are continuous families of foliations with non-trivial variations of their secondary classes for  $q \geq 3$ .

**Theorem:** (Rasmussen [1980]) There are continuous families of foliations with non-trivial variations of their secondary classes for  $q = 2$ .

**Corollary:**  $B\Gamma_q^r$  has uncountable topological type for all  $q \geq 1$ .

**Theorem:** (Hurder [1980]) For  $q \geq 2$ ,  $\pi_n(B\Gamma_q^r) \rightarrow \mathbb{R}^{k_n} \rightarrow 0$  where  $k_{2q+1} \neq 0$ , and in general,  $k_n$  has a subsequence  $k_{n_\ell} \rightarrow \infty$

# Homotopy chaos

**Theorem:** (Bott–Heitsch [1972])  $B\Gamma_q^r$  does not have finite topological type for  $q \geq 2$ .

**Theorem:** (Thurston [1972])  $\pi_3(B\Gamma_1^r)$  surjects onto  $\mathbb{R}$ .

**Theorem:** (Heitsch [1978]) There are continuous families of foliations with non-trivial variations of their secondary classes for  $q \geq 3$ .

**Theorem:** (Rasmussen [1980]) There are continuous families of foliations with non-trivial variations of their secondary classes for  $q = 2$ .

**Corollary:**  $B\Gamma_q^r$  has uncountable topological type for all  $q \geq 1$ .

**Theorem:** (Hurder [1980]) For  $q \geq 2$ ,  $\pi_n(B\Gamma_q^r) \rightarrow \mathbb{R}^{k_n} \rightarrow 0$  where  $k_{2q+1} \neq 0$ , and in general,  $k_n$  has a subsequence  $k_{n_\ell} \rightarrow \infty$

Secondary classes measure some uncountable aspect of foliation geometry.

## $C^2$ is essential

In contrast, Takashi Tsuboi proved the following amazing result:

**Theorem:** (Tsuboi [1989]) The classifying map of the normal bundle  $\nu: B\Gamma_q^1 \rightarrow BO(q)$  for foliations of transverse differentiability class  $C^1$  is a homotopy equivalence.

The proof is a technical tour-de-force, using Mather-Thurston type techniques for the study of  $B\Gamma$ .

## $C^2$ is essential

In contrast, Takashi Tsuboi proved the following amazing result:

**Theorem:** (Tsuboi [1989]) The classifying map of the normal bundle  $\nu: B\Gamma_q^1 \rightarrow BO(q)$  for foliations of transverse differentiability class  $C^1$  is a homotopy equivalence.

The proof is a technical tour-de-force, using Mather-Thurston type techniques for the study of  $B\Gamma$ .

When the  $C^1$  and  $C^2$  situations are radically different, one asks if there is some aspects of dynamical systems involved? (There are other reasons to ask this question, too.)

# Foliation dynamics

- A continuous dynamical system on a compact manifold  $M$  is a flow  $\varphi: M \times \mathbb{R} \rightarrow M$ , where the orbit  $L_x = \{\varphi_t(x) = \varphi(x, t) \mid t \in \mathbb{R}\}$  is thought of as the time trajectory of the point  $x \in M$ . The trajectories of the points of  $M$  are necessarily points, circles or lines immersed in  $M$ , and the study of their aggregate and statistical behavior is the subject of ergodic theory for flows.
- In foliation dynamics, we replace the concept of time-ordered trajectories with multi-dimensional futures for points. The study of the dynamics of  $\mathcal{F}$  asks for properties of the aggregate and statistical behavior of the collection of its leaves.



## Pseudogroups & Groupoids

Every foliation admits a discrete model by choosing a section  $\mathcal{T} \subset M$ , an embedded submanifold of dimension  $q$  which intersects each leaf of  $\mathcal{F}$  at least once, and always transversally. The holonomy of  $\mathcal{F}$  yields a compactly generated pseudogroup  $\mathcal{G}_{\mathcal{F}}$  acting on  $\mathcal{T}$ .

**Definition:** A pseudogroup of transformations  $\mathcal{G}$  of  $\mathcal{T}$  is *compactly generated* if there is

- a relatively compact open subset  $\mathcal{T}_0 \subset \mathcal{T}$  meeting all leaves of  $\mathcal{F}$
- a finite set  $\Gamma = \{g_1, \dots, g_k\} \subset \mathcal{G}$  such that  $\langle \Gamma \rangle = \mathcal{G}|_{\mathcal{T}_0}$ ;
- $g_i: D(g_i) \rightarrow R(g_i)$  is the restriction of  $\tilde{g}_i \in \mathcal{G}$  with  $\overline{D(g_i)} \subset D(\tilde{g}_i)$ .

## Pseudogroups & Groupoids

Every foliation admits a discrete model by choosing a section  $\mathcal{T} \subset M$ , an embedded submanifold of dimension  $q$  which intersects each leaf of  $\mathcal{F}$  at least once, and always transversally. The holonomy of  $\mathcal{F}$  yields a compactly generated pseudogroup  $\mathcal{G}_{\mathcal{F}}$  acting on  $\mathcal{T}$ .

**Definition:** A pseudogroup of transformations  $\mathcal{G}$  of  $\mathcal{T}$  is *compactly generated* if there is

- a relatively compact open subset  $\mathcal{T}_0 \subset \mathcal{T}$  meeting all leaves of  $\mathcal{F}$
- a finite set  $\Gamma = \{g_1, \dots, g_k\} \subset \mathcal{G}$  such that  $\langle \Gamma \rangle = \mathcal{G}|_{\mathcal{T}_0}$ ;
- $g_i: D(g_i) \rightarrow R(g_i)$  is the restriction of  $\tilde{g}_i \in \mathcal{G}$  with  $\overline{D(g)} \subset D(\tilde{g}_i)$ .

**Definition:** The groupoid of  $\mathcal{G}$  is the space of germs

$$\Gamma_{\mathcal{G}} = \{[g]_x \mid g \in \mathcal{G} \text{ \& } x \in D(g)\}, \quad \Gamma_{\mathcal{F}} = \Gamma_{\mathcal{G}_{\mathcal{F}}}$$

with source map  $s[g]_x = x$  and range map  $r[g]_x = g(x) = y$ .

## Derivative cocycle

Assume  $(\mathcal{G}, \mathcal{T})$  is a compactly generated pseudogroup, and  $\mathcal{T}$  has a uniform Riemannian metric. Choose a uniformly bounded, Borel trivialization,  $T\mathcal{T} \cong \mathcal{T} \times \mathbb{R}^q$ ,  $T_x\mathcal{T} \cong_x \mathbb{R}^q$  for all  $x \in \mathcal{T}$ .

**Definition:** The normal cocycle  $D\varphi: \Gamma_{\mathcal{G}} \times \mathcal{T} \rightarrow \mathbf{GL}(\mathbb{R}^q)$  is defined by

$$D\varphi[g]_x = D_x g: T_x\mathcal{T} \cong_x \mathbb{R}^q \rightarrow T_y\mathcal{T} \cong_y \mathbb{R}^q$$

which satisfies the cocycle law

$$D([h]_y \circ [g]_x) = D[h]_y \cdot D[g]_x$$

# Pseudogroup word length

**Definition:** For  $g \in \Gamma_{\mathcal{G}}$ , the word length  $\| [g] \|_x$  of the germ  $[g]_x$  of  $g$  at  $x$  is the least  $n$  such that

$$[g]_x = [g_{i_1}^{\pm 1} \circ \cdots \circ g_{i_n}^{\pm 1}]_x$$

Word length is a measure of the “time” required to get from one point on an orbit to another.

## Asymptotic exponent

**Definition:** The transverse expansion rate function at  $x$  is

$$\lambda(\mathcal{G}, n, x) = \max_{\|[\mathcal{G}]\|_x \leq n} \frac{\ln(\max\{\|D_x \mathcal{G}\|, \|D_y \mathcal{G}^{-1}\|\})}{\|[\mathcal{G}]\|_x} \geq 0$$

**Definition:** The asymptotic transverse growth rate at  $x$  is

$$\lambda(\mathcal{G}, x) = \limsup_{n \rightarrow \infty} \lambda(\mathcal{G}, n, x) \geq 0$$

This is essentially the maximum Lyapunov exponent for  $\mathcal{G}$  at  $x$ .

## Expansion classification

$$M = \mathcal{E} \cup \mathcal{P} \cup \mathcal{H}$$

where each are  $\mathcal{F}$ -saturated, Borel subsets of  $M$ , defined by:

# Expansion classification

$$M = \mathcal{E} \cup \mathcal{P} \cup \mathcal{H}$$

where each are  $\mathcal{F}$ -saturated, Borel subsets of  $M$ , defined by:

- 1 Elliptic points:  $\mathcal{E} \cap \mathcal{T} = \{x \in \mathcal{T} \mid \forall n \geq 0, \lambda(\mathcal{G}, n, x) \leq \kappa(x)\}$   
i.e., “points of bounded expansion” – e.g., Riemannian foliations

# Expansion classification

$$M = \mathcal{E} \cup \mathcal{P} \cup \mathcal{H}$$

where each are  $\mathcal{F}$ -saturated, Borel subsets of  $M$ , defined by:

- 1 Elliptic points:  $\mathcal{E} \cap \mathcal{T} = \{x \in \mathcal{T} \mid \forall n \geq 0, \lambda(\mathcal{G}, n, x) \leq \kappa(x)\}$   
i.e., “points of bounded expansion” – e.g., Riemannian foliations
- 2 Parabolic points:  $\mathcal{P} \cap \mathcal{T} = \{x \in \mathcal{T} - (\mathcal{E} \cap \mathcal{T}) \mid \lambda(\mathcal{G}, x) = 0\}$   
i.e., “points of slow-growth expansion” – e.g., distal foliations



# Expansion classification

$$M = \mathcal{E} \cup \mathcal{P} \cup \mathcal{H}$$

where each are  $\mathcal{F}$ -saturated, Borel subsets of  $M$ , defined by:

- 1 Elliptic points:  $\mathcal{E} \cap \mathcal{T} = \{x \in \mathcal{T} \mid \forall n \geq 0, \lambda(\mathcal{G}, n, x) \leq \kappa(x)\}$   
i.e., “points of bounded expansion” – e.g., Riemannian foliations
- 2 Parabolic points:  $\mathcal{P} \cap \mathcal{T} = \{x \in \mathcal{T} - (\mathcal{E} \cap \mathcal{T}) \mid \lambda(\mathcal{G}, x) = 0\}$   
i.e., “points of slow-growth expansion” – e.g., distal foliations
- 3 Partially Hyperbolic points:  $\mathcal{H} \cap \mathcal{T} = \{x \in \mathcal{T} \mid \lambda(\mathcal{G}, x) > 0\}$   
i.e., “points of exponential-growth expansion” – non-uniformly, partially hyperbolic foliations

## Secondary classes and dynamics

A secondary class  $h_I c_J \in H^*(WO_q)$  is *residual* if  $c_J$  has degree  $2q$ .

**Theorem:** (Hurder, 2006) Let  $h_I c_J \in H^*(WO_q)$  be a residual secondary class (e.g., Godbillon-Vey type). Suppose that  $\Delta_{\mathcal{F}}(h_I c_J) \in H^*(M; \mathbb{R})$  is non-zero. Then the hyperbolic component  $\mathcal{H}$  has positive Lebesgue measure.

Moreover, the elliptic  $\mathcal{E}$  and parabolic  $\mathcal{P}$  components do not contribute to the secondary classes. (i.e., The Weil measure for  $h_I$  vanishes on these components, hence the restrictions of the residual secondary classes to these sets are trivial in cohomology.)

## Secondary classes and dynamics

A secondary class  $h_{Ic_J} \in H^*(WO_q)$  is *residual* if  $c_J$  has degree  $2q$ .

**Theorem:** (Hurder, 2006) Let  $h_{Ic_J} \in H^*(WO_q)$  be a residual secondary class (e.g., Godbillon-Vey type). Suppose that  $\Delta_{\mathcal{F}}(h_{Ic_J}) \in H^*(M; \mathbb{R})$  is non-zero. Then the hyperbolic component  $\mathcal{H}$  has positive Lebesgue measure.

Moreover, the elliptic  $\mathcal{E}$  and parabolic  $\mathcal{P}$  components do not contribute to the secondary classes. (i.e., The Weil measure for  $h_I$  vanishes on these components, hence the restrictions of the residual secondary classes to these sets are trivial in cohomology.)

Understanding the “dynamical meaning of the residual secondary classes” in  $H^*(WO_q)$  requires understanding the dynamics of foliations which have non-uniformly, partially hyperbolic behavior on a set of positive measure.

# Framed foliations

But... is this a true picture of the relation between topology and dynamics?

## Framed foliations

But... is this a true picture of the relation between topology and dynamics?

**Definition:**  $\mathcal{F}$  is framed if there is a framing  $s: M \rightarrow \mathbf{Fr}(Q)$  of the normal bundle  $Q \rightarrow M$ . The classifying space  $F\Gamma_q$  of framed foliations is the homotopy fiber

$$F\Gamma_q \rightarrow B\Gamma_q \rightarrow BO_q$$

## Framed foliations

But... is this a true picture of the relation between topology and dynamics?

**Definition:**  $\mathcal{F}$  is framed if there is a framing  $s: M \rightarrow \mathbf{Fr}(Q)$  of the normal bundle  $Q \rightarrow M$ . The classifying space  $F\Gamma_q$  of framed foliations is the homotopy fiber

$$F\Gamma_q \rightarrow B\Gamma_q \rightarrow BO_q$$

The transgressions of the Pontrjagin classes  $p_j = c_{2j}$  are now defined:

$$W_q \cong \Lambda(h_1, h_2, \dots, h_{q/2}) \otimes \mathbb{R}[c_1, c_2, \dots, c_q], \quad d_W h_i = c_i, \quad d_W c_i = 0$$

**Theorem':** (Bott-Haefliger, Gelfand-Fuks, Kamber-Tondeur [1972])

There is a functorial characteristic map

$$\Delta^s: H^*(W_q) \rightarrow H^*(F\Gamma_q; \mathbb{R})$$

Classes involving the terms  $h_{2i}$  can also vary in examples.

# Minimal sets

Introduce another basic idea of dynamics:

**Definition:** A closed, saturated subset  $K \subset M$  is *minimal* if every leaf  $L \subset K$  is dense in  $K$ .

# Minimal sets

Introduce another basic idea of dynamics:

**Definition:** A closed, saturated subset  $K \subset M$  is *minimal* if every leaf  $L \subset K$  is dense in  $K$ .

A minimal set  $K$  can be one of three types:

- $K = L$  is a compact leaf of  $\mathcal{F}$
- $K$  has interior, hence  $M$  connected implies  $K = M$
- $K$  is not a leaf, and has no interior, hence  $K$  is a perfect subset.

The latter case is called an *exceptional* minimal set for historical reasons.



# An essential exceptional parabolic minimal set

**Theorem:** (Hurder, 2008) For  $q \geq 2$ , there exists a framed foliation  $\mathcal{F}$  with exceptional minimal set  $\mathcal{S}$  such that:

- $\mathcal{F}$  is a parabolic foliation –  $\mathcal{S}$  has no transverse hyperbolicity
- For every open neighborhood  $\mathcal{S} \subset U$ , the classifying map  $h_{\mathcal{F}}: U \rightarrow F\Gamma_q$  is not homotopically trivial.

# An essential exceptional parabolic minimal set

**Theorem:** (Hurder, 2008) For  $q \geq 2$ , there exists a framed foliation  $\mathcal{F}$  with exceptional minimal set  $\mathcal{S}$  such that:

- $\mathcal{F}$  is a parabolic foliation –  $\mathcal{S}$  has no transverse hyperbolicity
- For every open neighborhood  $\mathcal{S} \subset U$ , the classifying map  $h_{\mathcal{F}}: U \rightarrow F\Gamma_q$  is not homotopically trivial.

$\mathcal{S}$  is a generalized solenoid, which is transversally a Cantor set  $\mathcal{C}$ , and the holonomy of  $\mathcal{F}$  restricted to  $\mathcal{C}$  is equivalent to an “adding machine”.

## Bott-Heitsch revisited

For the construction of  $\mathcal{S}$ , we go back to the beginning:

**Theorem:** (Bott-Heitsch [1972])

$h_Q^*: H^*(BO_q; \mathbb{Z}) \rightarrow H^*(M; \mathbb{Z})$  is injective for all  $*$ .

## Bott-Heitsch revisited

For the construction of  $\mathcal{S}$ , we go back to the beginning:

**Theorem:** (Bott-Heitsch [1972])

$h_Q^*: H^*(BO_q; \mathbb{Z}) \rightarrow H^*(M; \mathbb{Z})$  is injective for all  $*$ .

We recall the proof for the case of oriented normal bundles and  $q = 2$ .

## Bott-Heitsch revisited

For the construction of  $\mathcal{S}$ , we go back to the beginning:

**Theorem:** (Bott-Heitsch [1972])

$h_Q^*: H^*(BO_q; \mathbb{Z}) \rightarrow H^*(M; \mathbb{Z})$  is injective for all  $*$ .

We recall the proof for the case of oriented normal bundles and  $q = 2$ .

$$H^*(BSO_2; \mathbb{Z}) \cong \mathbb{Z}[e]$$

Let  $n > 2$ , and set  $\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$ . Embed  $\mathbb{Z}_n \subset SO_2$ , acts isometrically on  $\mathbb{R}^2$   
 $\mathbb{Z}_n$  acts freely on  $\mathbb{S}^{2k+1}$  for  $k > 0$ .

$$\mathbb{E}_{n,k} = \mathbb{S}^{2k} \times \mathbb{R}^2 / \mathbb{Z}_n, \quad \mathcal{F}_{n,k} = \text{flat bundle foliation}$$

For  $* \leq 2k$  have injection:

$$\mathbb{Z}_n[e] \rightarrow H^*(BSO_2; \mathbb{Z}_n) \rightarrow H^*(B\Gamma_2; \mathbb{Z}_n) \rightarrow H^*(\mathbb{E}_{n,\ell}; \mathbb{Z}_n)$$

So  $H^*(BSO_2; \mathbb{Z}_n) \rightarrow H^*(B\Gamma_2; \mathbb{Z}_n)$  is injective for all  $*$  and all  $n \rightarrow \infty$ .

$H^*(BSO_2; \mathbb{Z}) \rightarrow H^*(B\Gamma_2; \mathbb{Z})$  injective follows from this.

For  $* \leq 2k$  have injection:

$$\mathbb{Z}_n[e] \rightarrow H^*(BSO_2; \mathbb{Z}_n) \rightarrow H^*(B\Gamma_2; \mathbb{Z}_n) \rightarrow H^*(\mathbb{E}_{n,\ell}; \mathbb{Z}_n)$$

So  $H^*(BSO_2; \mathbb{Z}_n) \rightarrow H^*(B\Gamma_2; \mathbb{Z}_n)$  is injective for all  $*$  and all  $n \rightarrow \infty$ .

$H^*(BSO_2; \mathbb{Z}) \rightarrow H^*(B\Gamma_2; \mathbb{Z})$  injective follows from this.

General case for  $q > 2$  uses splitting principle, for torsion subgroups of maximal torus,  $\mathbb{Z}_n^k \subset \mathbb{T}^k \subset SO_{2k}$

For  $* \leq 2k$  have injection:

$$\mathbb{Z}_n[e] \rightarrow H^*(BSO_2; \mathbb{Z}_n) \rightarrow H^*(B\Gamma_2; \mathbb{Z}_n) \rightarrow H^*(\mathbb{E}_{n,\ell}; \mathbb{Z}_n)$$

So  $H^*(BSO_2; \mathbb{Z}_n) \rightarrow H^*(B\Gamma_2; \mathbb{Z}_n)$  is injective for all  $*$  and all  $n \rightarrow \infty$ .

$H^*(BSO_2; \mathbb{Z}) \rightarrow H^*(B\Gamma_2; \mathbb{Z})$  injective follows from this.

General case for  $q > 2$  uses splitting principle, for torsion subgroups of maximal torus,  $\mathbb{Z}_n^k \subset \mathbb{T}^k \subset SO_{2k}$

**Question:** Can we realize this limit process  $(n, k) \rightarrow \infty$  with foliation?




## Dynamics of flat bundles

Switch to groupoid model:  $\mathbb{Z}_n$  acting on disk  $\mathbb{D}^2 \subset \mathbb{R}^2$  via rotations.

Action is free except at center point of disk.

Pick  $0 \neq z_1 \in \mathbb{D}^2$ , with orbit  $\mathbb{Z}_n \cdot z_1 = \{z_{1,0}, \dots, z_{1,n-1}\}$ .

Consider disks  $\mathbb{D}_{1,i}^2(z_{1,i}, \epsilon_1) \subset \mathbb{D}^2$  for  $\epsilon_1 > 0$  sufficiently small. Here is illustration in case of  $n = 6$ :



# Semi-simplicial realization of flat bundles

Let  $\Gamma_{2,n} = (\mathbb{D}^2, \mathbb{Z}_n)$  denote the associated groupoid.

$|\Gamma_{2,n}|$  is the semi-simplicial space realizing the groupoid.

Then the classifying map factors:

$$\mathbb{E}_{n,k} \rightarrow |\Gamma_{2,n}| \rightarrow B\Gamma_2$$

**Corollary:**  $H^*(BSO_2; \mathbb{Z}_n) \rightarrow H^*(B\Gamma_2; \mathbb{Z}_n) \rightarrow H^*(|\Gamma_{2,n}|; \mathbb{Z}_n)$   
is injective for all  $*$  and all  $n \rightarrow \infty$ .

## Construction of solenoids

Choose  $n_1 < n_2 < \dots$  tending to infinity rapidly. Example:  $n_k = 3^{k!}$

Choose  $\epsilon_k \rightarrow 0$  rapidly, but slower than  $1/n_k$ . Example:  $\epsilon_n = \epsilon_0 \cdot (3^n d_n)^{-1}$

Restriction of  $\Gamma_{2,n_1} = (\mathbb{D}^2, \mathbb{Z}_{n_1})$  to the invariant set

$$\mathcal{S}_1 = \mathbb{D}_{1,0}^2(z_{1,0}, \epsilon_1) \cup \mathbb{D}_{1,1}^2(z_{1,1}, \epsilon_1) \cup \dots \cup \mathbb{D}_{n-1}^2(z_{1,n_1-1}, \epsilon_1)$$

is free, so we can repeat this construction of an action on  $\mathcal{S}_1$ .


Choose  $z_{2,0} \in \mathbb{D}_{1,0}^2(z_{1,0}, \epsilon_1)$  which is not on center.

Repeat above construction for  $\mathbb{Z}_{n_2}$  on disks  $\mathbb{D}_{2,0}^2(z_{2,0}, \epsilon_2)$ .

There is one catch! Cannot just insert this action into  $\Gamma_{2,n_1}$ . The plug will not be smooth.

Need to make deformation of action from identity on boundary of  $V_1$  to rotation by  $2\pi/n_2$  on boundary of  $\mathbb{D}_{2,0}^2(z_{2,0}, \epsilon_2)$ .

# Picture of stage 1: $n_1 = 2$



## Limit solenoid

Let  $\Gamma_{2,\infty}$  the smooth groupoid resulting from the limit of this construction.  
The action on  $\mathbb{D}^2$  is distal!

**Proposition:** The dynamics of  $\Gamma_{2,\infty}$  contains a solenoidal minimal set

$$\mathcal{S} = \bigcap_{k=1}^{\infty} |\mathcal{S}_k|$$

**Proposition:** For every open neighborhood  $\mathcal{S} \subset U \subset |\Gamma_{2,\infty}|$  there exists some  $k \gg 0$  such that  $|\mathcal{S}_k| \subset U$

**Corollary:** For  $k \gg 0$  there is an inclusion  $|\Gamma_{2,k}| \subset |\Gamma_{2,\infty}|$ .

## Homotopical consequences

Let  $U$  be an open neighborhood,  $\mathcal{S} \subset U \subset |\Gamma_{2,\infty}|$ .

**Proposition:**  $H^*(BSO_2; \mathbb{Z}) \rightarrow H^*(B\Gamma_2; \mathbb{Z}) \rightarrow H^*(U; \mathbb{Z})$  is injective.

## Homotopical consequences

Let  $U$  be an open neighborhood,  $\mathcal{S} \subset U \subset |\Gamma_{2,\infty}|$ .

**Proposition:**  $H^*(BSO_2; \mathbb{Z}) \rightarrow H^*(B\Gamma_2; \mathbb{Z}) \rightarrow H^*(U; \mathbb{Z})$  is injective.

**Corollary:** The image of the classifying map  $U \rightarrow B\Gamma_2$  cannot have finite type in all odd dimensions  $> 4$ .

## Homotopical consequences

Let  $U$  be an open neighborhood,  $\mathcal{S} \subset U \subset |\Gamma_{2,\infty}|$ .

**Proposition:**  $H^*(BSO_2; \mathbb{Z}) \rightarrow H^*(B\Gamma_2; \mathbb{Z}) \rightarrow H^*(U; \mathbb{Z})$  is injective.

**Corollary:** The image of the classifying map  $U \rightarrow B\Gamma_2$  cannot have finite type in all odd dimensions  $> 4$ .

One obtains framed foliations by considering the frame bundle  $\widehat{U} \rightarrow U$  of the normal bundle on  $U$ .

The foliation  $\mathcal{F}$  on  $U$  lifts to a foliation  $\widehat{\mathcal{F}}$  on  $\widehat{U}$ .

By finite-type considerations, we obtain

**Theorem:** The image of the classifying map  $\widehat{U} \rightarrow F\Gamma_q$  cannot have finite type in all odd dimensions  $> 4$ .



# Chern-Simons invariants

**Theorem:** The Chern-Simons invariants in  $H^{2^*-1}(B\Gamma_2; \mathbb{R}/\mathbb{Z})$  are non-trivial on the image of  $|\Gamma_{2,\infty}| \rightarrow B\Gamma_q$  in all odd dimensions  $> 4$ .

# Chern-Simons invariants

**Theorem:** The Chern-Simons invariants in  $H^{2*-1}(B\Gamma_2; \mathbb{R}/\mathbb{Z})$  are non-trivial on the image of  $|\Gamma_{2,\infty}| \rightarrow B\Gamma_q$  in all odd dimensions  $> 4$ .

**Remark 1:** Apparently, the transgression classes of the Pontrjagin classes  $H^{4*}(BSO_q; \mathbb{R})$  do not depend on dynamics in the same way as before.

# Chern-Simons invariants

**Theorem:** The Chern-Simons invariants in  $H^{2*-1}(B\Gamma_2; \mathbb{R}/\mathbb{Z})$  are non-trivial on the image of  $|\Gamma_{2,\infty}| \rightarrow B\Gamma_q$  in all odd dimensions  $> 4$ .

**Remark 1:** Apparently, the transgression classes of the Pontrjagin classes  $H^{4*}(BSO_q; \mathbb{R})$  do not depend on dynamics in the same way as before.

**Remark 2:** The above construction admits many generalizations to embedded braid diagrams. Unclear what cohomology theories will be needed to detect them.